

An elementary illustrated introduction to simplicial sets

Greg Friedman
Texas Christian University

September 23, 2008

2000 Mathematics Subject Classification: 18G30, 55U10

Keywords: Simplicial sets, simplicial homotopy

Abstract

This is an expository introduction to simplicial sets and simplicial homotopy theory with particular focus on relating the combinatorial aspects of the theory to their geometric/topological origins. It is intended to be accessible to students familiar with just the fundamentals of algebraic topology.

Contents

1	Introduction	2
2	A build-up to simplicial sets	3
2.1	Simplicial complexes	3
2.2	Simplicial maps	6
2.3	Delta sets and Delta maps	7
3	Simplicial sets and morphisms	11
4	Realization	19
5	Products	24
6	Simplicial objects in other categories	28
7	Kan complexes	30
8	Simplicial homotopy	32
8.1	Paths and path components	32
8.2	Homotopies of maps	34
8.3	Relative homotopy	36

1 Introduction

The following notes grew out of my own difficulties in attempting to learn the basics of simplicial sets and simplicial homotopy theory, and thus they are aimed at someone with roughly the same starting knowledge I had, specifically some amount of comfort with simplicial homology (e.g. that available to those of us who grew up learning homology from Munkres [14], slightly less so for those coming of age in Hatcher [9]) and the basic fundamentals of topological homotopy theory, including homotopy groups. Equipped with this background, I wanted to understand a little of what simplicial sets and their generalizations to other categories are all about, as they seem ubiquitous in the literature of certain schools of topology. To name just a few important instances of which I am aware, simplicial objects occur in May's work on recognition principles for iterated loop spaces [10], Quillen's approach to rational homotopy theory (see [16, 6]), Bousfield and Kan's work on completions, localization, and limits in homotopy theory [1], Quillen's abstract treatment of homotopy theory [17], and various aspects of homological algebra, including group cohomology, Hochschild homology, and cyclic homology (see [21]).

However, in attempting to learn the rudiments of simplicial theory, I encountered immediate and discouraging difficulties, which led to serious frustration on several occasions. It was only after several different attempts from different angles that I finally began to "see the picture," and my intended goal here is to aid future students (of all ages) to ease into the subject.

My initial difficulty with the classic expository sources such as May [11] and Curtis [3] was the extent to which the theory is presented purely combinatorially. And the combinatorial definitions are not often pretty; they tend to consist of long strings of axiomatic conditions (see, for example, the combinatorial definition of simplicial homotopy, Definition 8.6, below). Despite simplicial objects originating in very topological settings, these classic expositions often sweep this fact too far under the rug for my taste, as someone who likes to comprehend even algebraic and combinatorial constructions as visually as possible. There is a little bit more geometry in Moore's lecture notes [13], though still not much, and these are also more difficult to obtain (at least not without some good help from a solid Interlibrary Loan Department). On the other hand, there is a much more modern point of view that sweeps both topology and combinatorics away in favor of axiomatic category theory! Goerss and Jardine [8] is an excellent modern text based upon this approach, which, ironically, helped me tremendously to understand what the combinatorics were getting at!

So what are we getting at here? My goal, still as someone very far from an expert in either combinatorial or axiomatic simplicial theory, is to revisit the material covered in, roughly, the first chapters (in some cases the first few pages) of the texts cited above and to provide some concrete geometric signposts. Here, for the most part, you won't find many complete proofs of theorems, and so these notes will not be completely self-contained. Rather, I try

primarily to show by example how the very basic combinatorics, including the definitions, arise out of geometric ideas and to show the geometric ideas underlying the most elementary proofs and properties. Think of this as an appendix or a set of footnotes to the first chapters of the classic expositions, or perhaps as a Chapter 0. This may not sound like much, but during my earliest learning stages with this material, I would have been very grateful for something of the sort. Theoretically my reader will acquire enough of “the idea” to go forth and read the more thorough (and more technical) sources equipped with enough intuition to *see* what’s going on.

In Section 2, we lay the groundwork with a look at the more familiar topics of simplicial sets and, their slight generalizations, Delta sets. Simplicial sets are then introduced in Section 3, followed by their geometric realizations in Section 4 and a detailed look at products of simplicial sets in Section 5. In Section 6, we provide a brief look at how the notion of simplicial sets is generalized to other kinds of simplicial objects based in different categories. In Section 7, we introduce Kan complexes; these are the simplicial sets that lend themselves to simplicial analogues of homotopy theory, which we study in Section 8. This section gets a bit more technical as we head toward more serious applications and theorems in simplicial theory, including the definition and properties of the simplicial homotopy groups $\pi_n(X, *)$ in Section 9. Finally, in Section 10, we make some concluding remarks and steer the reader toward more comprehensive expository sources.

2 A build-up to simplicial sets

We begin at the beginning with the relevant geometric notions and their immediate combinatorial counterparts.

2.1 Simplicial complexes

Simplicial objects are, essentially, generalizations of the geometric simplicial complexes of elementary algebraic topology (in some cases quite extreme generalizations). So let’s recall simplicial complexes, referring the absolute beginner to [14] for a complete course in the essentials.

Recall that a simplicial complex X is made up of simplices (generalized tetrahedra) of various dimensions, glued together along common faces (see Figure 1). The most efficient description, containing all of the relevant information, comes from labeling the vertices (the 0-simplices) and then specifying which collections of vertices together constitute the vertices of simplices of higher dimension. Most often one assumes that the collection of vertices is countable so that we can label them $v_0, v_1, v_2, \dots$, though this is not strictly necessary - we could label by elements of any set. Then if some collection of vertices $\{v_{i_0}, \dots, v_{i_n}\}$ constitutes the vertices of a simplex, we can write that simplex as $[v_{i_0}, \dots, v_{i_n}]$.

A nice way to organize the combinatorial information involved is to define the skeleta X^k , $k = 0, 1, \dots$, of a simplicial complex so that X^k is the set of all k -simplices of X . Notice that, having labeled our vertices so that $X^0 = \{v_i\}_{i=0}^\infty$, we can think of the elements of X^k




Figure 1: A simplicial complex. Note that $[v_0, v_1, v_2]$ is a simplex, but $[v_1, v_2, v_3]$ is not.

as subsets of X^0 ; a subset $\{v_{i_0}, \dots, v_{i_k}\} \subset X^0$ is an *element* of X^k precisely if $[v_{i_0}, \dots, v_{i_k}]$ is a k -dimensional simplex of X .

If X is a complex and $[v_{i_0}, \dots, v_{i_k}]$ is a simplex of X , then any subset of $\{v_{i_0}, \dots, v_{i_k}\}$ is a face of that simplex and thus itself a simplex of X . This observation and this type of notation leads us directly to the notion of an abstract simplicial complex:

Definition 2.1. An *abstract simplicial complex* consists of a set of “vertices” X^0 together with, for each integer k , a set X^k consisting of subsets of X^0 of cardinality $k + 1$. These must satisfy the condition that any $j + 1$ -element subset of an element of X^k is an element of X^j .

So, an abstract simplicial complex has exactly the same combinatorial information as a geometric simplicial complex (what we have lost is geometric information about how big a simplex is, what are its dimensions, how is it embedded in euclidean space, etc.), but we have retained all of the information necessary to reconstruct the complex up to homeomorphism. It is straightforward that a geometric complex yields an abstract complex, but conversely, we can obtain a geometric complex (up to homeomorphism) from an abstract one by assigning to each element of X^0 a geometric point and to each abstract simplex $[v_{i_0}, \dots, v_{i_k}]$ a geometric k -simplex spanned by the appropriate vertices and gluing these simplices together via the quotient topology.

Ordered simplicial complexes. A slightly more specific way to do all this is to let our set of vertices X^0 be totally ordered, in which case we obtain an ordered simplicial complex. When we do this, $[v_{i_0}, \dots, v_{i_k}]$ may stand for a simplex if and only if $v_{i_j} < v_{i_l}$ whenever $j < l$. This poses no undue complications as each collection $\{v_{i_0}, \dots, v_{i_k}\}$ of cardinality k still corresponds to at most one simplex. We’re just being picky and removing some redundancy in how many ways we can label a given simplex of a simplicial complex.

Example 2.2. Of course the prototypical example of a simplicial complex is the n -simplex itself, denoted $|\Delta^n|$ (see Figure 2); it will become clear later why we want to employ the notation $|\Delta^n|$ instead of just Δ^n .




Figure 2: The standard 0-, 1-, 2-, and 3-simplices

The n -simplex is so fundamental that one often labels the vertices simply with the numbers $0, 1, \dots, n$. This notation should be suggestive when compared with the simplices $[v_{i_0}, \dots, v_{i_n}]$ appearing within more general simplicial complexes, and it is worth pointing out at this early stage that one can think of any such simplex in a complex X as the image of $|\Delta^n|$ under a simplicial map taking 0 to v_{i_0} , and so on; we will discuss simplicial maps more formally in a moment. If X is an ordered complex, then there is precisely one way to do this for each simplex of X . Thus another point of view on ordered simplicial complexes is that they are made up out of images of the standard simplices (Figure 3). This will turn out to be a very useful point of view as we progress.




Figure 3: $[v_2, v_3, v_4]$ as the image of $|\Delta^2|$

Note also that the faces of the standard n -simplex can be represented simply as collections of numbers $[j_1, \dots, j_m]$ with $0 \leq j_1 < j_2 < \dots < j_m \leq n$ and, as an ordered simplicial complex, the set of simplices (of all dimensions) in $|\Delta^n|$ is equivalent to the collection of ordered subsets of $\{0, \dots, n\}$.

Face maps. Another aspect of ordered simplicial complexes familiar to the student of basic algebraic topology is that, given an n -simplex, we would like a handy way of referring to its $n - 1$ faces. This is handled by the *face maps*. On the standard n -simplex, we have $n + 1$ face maps $d_0, \dots, d_n$, defined so that $d_j[0, \dots, n] = [0, \dots, \hat{j}, \dots, n]$, where, as usual, the $\hat{}$ denotes a term that is being omitted. Thus applying d_j to $[0, \dots, n]$ picks out the $n - 1$ face missing the vertex j . (See Figure 4).

Within more general simplicial complexes, we make the obvious extension: if $[v_{i_0}, \dots, v_{i_n}] \in X^n$ is a simplex of the complex X , then $d_j[v_{i_0}, \dots, v_{i_n}] = [v_{i_0}, \dots, \hat{v}_{i_j}, \dots, v_{i_n}]$. Assembled




Figure 4: The face maps of $|\Delta^2|$

all together, we get, for each fixed n , a collection of functions $d_0, \dots, d_n : X^n \rightarrow X^{n-1}$. Note that here is where the ordering of the vertices of the simplices becomes important.

If one wanted to be a serious stickler, we might be careful to label the face maps from X^n to X^{n-1} as $d_0^n, \dots, d_n^n$, but this is rarely done in practice, for which we should probably be grateful. Thus d_j is used to represent the face map leaving out the j th vertex in any dimension where this makes sense (i.e. dimensions $\geq j$).

Furthermore, one readily sees by playing with $|\Delta^n|$ that there are certain relations satisfied by the face maps. In particular, if $i < j$, then $d_i d_j = d_{j-1} d_i$. Indeed, $d_i d_j[0, \dots, n] = [0, \dots, \hat{i}, \dots, \hat{j}, \dots, n] = d_{j-1} d_i[0, \dots, n]$ (notice the reason that we have d_{j-1} is the last expression is that removing the i first shifts the j into the $j-1$ slot).

Clearly, the relation $d_i d_j = d_{j-1} d_i$ must hold for any simplex in a complex X (which is made up of copies of $|\Delta^n|$). This relation will become one of the axioms in the definition of a simplicial set when we get there.

Another observation that will come up later is that there are more general face maps. We could, for example, assign to $[0, 1, 2, 3, 4, 5, 6]$ the face $[1, 3, 4]$, and we could define such general face maps systematically. However, any such face can be obtained as a combination of face maps that lower dimension by 1. For example, we can decompose the map just described as $d_0 d_2 d_5 d_6$. It may entertain the reader to use the “face map relations” and some basic reasoning to show that any generalized face map can be obtained as a composition $d_{i_1} \cdots d_{i_m}$ *uniquely* if we require that $i_j < i_{j+1}$ for all j .

2.2 Simplicial maps

The heart of the transition (in my opinion) from simplicial complexes to simplicial sets is tied up in simplicial maps.

Recall (see [14, Section 2]) that if K and L are simplicial complexes, then a simplicial map $f : K \rightarrow L$ is determined by taking the vertices $\{v_i\}$ to vertices $\{f(v_i)\}$ of L such that if $[v_{i_0}, \dots, v_{i_k}]$ is a simplex of K then $f(v_{i_0}), \dots, f(v_{i_k})$ are all vertices (not necessarily unique) of some simplex in L . Given such a function $K^0 \rightarrow L^0$, the rest of $f : K \rightarrow L$ is determined

by linear interpolation on each simplex (if $x \in K$ can be represented by $x = \sum_{j=1}^n t_j v_{i_j}$ in barycentric coordinates of the simplex spanned by the v_{i_j} , then $f(x) = \sum_{j=1}^n t_j f(v_{i_j})$). The resulting function $f : K \rightarrow L$ is continuous (see [14]).

Example 2.3. We have already seen one example of a simplicial map in Example 2.2, the one that takes the n -simplex $|\Delta^n|$ to a simplex $[v_{i_0}, \dots, v_{i_n}]$ of some simplicial complex.

Example 2.4. The interesting feature of simplicial maps, from the point of view of simplicial sets, is that simplicial maps can collapse simplices. For example, consider the simplicial map $f : \Delta^2 \rightarrow \Delta^1$ determined by $f(0) = 0$, $f(1) = 1$, $f(2) = 1$ that collapses the 2-simplex down to the 1-simplex (see Figure 5). The great benefit of the theory of simplicial sets is a way to generalize these kinds of maps in order to preserve information so that we can still see the image of the 2-simplex hiding in the 1-simplex as a *degenerate* simplex (see Section 3).




Figure 5: A collapse of $|\Delta^2|$ to $|\Delta^1|$

2.3 Delta sets and Delta maps

Delta sets (sometimes called Δ -sets) constitute an intermediary between simplicial complexes and simplicial sets. These allow a degree of abstraction without yet introducing the degeneracy maps we have begun hinting at.

Definition 2.5. A *Delta set*¹ consists of a sequence of ordered sets $X_0, X_1, \dots$ and, for each $n \geq 0$, maps $d_i : X_{n+1} \rightarrow X_n$ for each i , $0 \leq i \leq n+1$, such that $d_i d_j = d_{j-1} d_i$ whenever $i < j$.

Of course this is just an abstraction, and generalization, of the definition of an ordered simplicial complex, in which the X_n are the sets of n -simplices and the d_i are the face maps. However, there are Delta sets that are not simplicial complexes:

¹It seems to be at least fairly usual to capitalize the word “Delta” in this context, probably since it is essentially a stand-in for the Greek capital letter Δ . However, for reasons that will become clear, it is probably best to avoid the notation “ Δ -set” and to use instead the English stand-in.

Example 2.6. Consider the cone C obtained by starting with the standard 2-simplex $|\Delta^2| = [0, 1, 2]$ and gluing the edge $[0, 2]$ to the edge $[1, 2]$ (see Figure 6). This space is no longer a simplicial complex (at least not with the “triangulation” given), since in a simplicial complex, the faces of a given simplex must be unique. This is no longer the case here as, for example, the “edge $[0, 1]$ ” now has both endpoint vertices equal to each other.



Figure 6: Gluing $|\Delta^2|$ into a cone

However, this is a Delta set. Without (I hope!) too much risk of confusion, we use the notation for the simplices in the triangle to refer also to their images in the cone. So, for example $[0]$ and $[1]$ now both stand for the same vertex in the cone and $[0, 1]$ stands for the circular base edge. Then $C_0 = \{[0], [2]\}$, $C_1 = \{[0, 1], [0, 2]\}$, $C_2 = [0, 1, 2]$, and $C_n = \emptyset$ for all $n > 2$. The face maps are the obvious ones, also induced from the triangle, so that, e.g. $d_2[0, 1, 2] = [0, 1]$ and $d_0[0, 1] = d_1[0, 1] = [0] = [1]$. It is not hard to see that the axiom on the d_i is satisfied - it comes right from the fact that it holds for the standard 2-simplex.

Example 2.7. One feature of Delta sets we need to be careful about is that, unlike for simplicial complexes, a collection of vertices does not necessarily specify a unique simplex. For example, consider the Delta set with $X_0 = \{v_0, v_1\}$, $X_1 = \{e_0, e_1\}$, $d_0(e_0) = d_0(e_1) = v_0$, and $d_1(e_0) = d_1(e_1) = v_1$. Both 1-simplices have the same endpoints. See Figure 7.




Figure 7: A Delta set containing two edges with the same vertices

Thus Delta sets afford some greater flexibility beyond ordered simplicial complexes. One may continue to think of the sets X_n as collections of simplices and interpret from the face maps how these are meant to be glued together (Exercise: Give each “simplex” of the cone X of the preceding example an abstract label, write out the full set of face maps in these labels, then reverse engineer how to construct the cone from this information. One sees that everything is forced. For example, there is one 2-simplex, two of whose faces are the same, so they must be glued together!). However, it is common in the fancier literature not to think of the X_n as collections of simplices at all but simply as abstract sets with abstract collections of face maps. At least this is what authors would have us believe - I tend to picture simplices in my head anyway, while keeping in mind that this is more of a cognitive aid than it is “what’s really going on.”

The category-theoretic definition. While we're walking the tightrope of abstraction, let's take it a step further. Recall that we discussed in Example 2.2 that we can think of an ordered simplicial complex as a collection of isomorphic images of the standard n -simplices. Of course to describe the simplicial complex fully we need not only the images of the standard simplices but we need to know which image faces are attached together. This information is contained in the face maps, which tell us when two simplices share a face. There's an alternative definition of Delta complexes that takes more of this point of view. It might be a little scary if you're not that comfortable with category theory, but don't worry I'll walk you through it (though I do assume you know the basic language of categories and functors).

First, we define a category $\hat{\Delta}$, $\hat{\Delta}$:

Definition 2.8. The category $\hat{\Delta}$ has as objects the finite ordered sets $[n] = \{0, 1, 2, \dots, n\}$. The morphisms of $\hat{\Delta}$ are the strictly order preserving functions $[m] \rightarrow [n]$ (recall that f is strictly order preserving if $i < j$ implies $f(i) < f(j)$).

Of course the objects of $\hat{\Delta}$ should be thought of as our prototype *ordered* n -simplices. The morphisms are only defined when $m \leq n$, and you can think of these morphisms as taking an m -simplex and embedding it as a face of an n -simplex (see figure 8. Note that, since order matters, there are exactly as many ways to do this as there are strictly order preserving maps $[m] \rightarrow [n]$).




Figure 8: A partial illustration of the category $\hat{\Delta}$

Next, we think about the opposite category $\hat{\Delta}^{op}$. Recall that this means that we keep the same objects $[n]$ of $\hat{\Delta}$, but for every morphism $[m] \rightarrow [n]$ in $\hat{\Delta}$, we instead have a map $[n] \rightarrow [m]$ in $\hat{\Delta}^{op}$. What should this mean? Well if a given morphism $[m] \rightarrow [n]$ was the

inclusion of a face, then the new opposite map $[n] \rightarrow [m]$ should be thought of as taking the n -simplex $[n]$ and prescribing a given face. This is just a generalization of what we have seen already: if we consider in $\widehat{\Delta}$ the morphism $D_i : [n] \rightarrow [n+1]$ defined by the strictly order preserving map $\{0, \dots, n\} \rightarrow \{0, \dots, \hat{i}, \dots, n+1\}$, then in $\widehat{\Delta}^{op}$ this corresponds precisely to the simplex face map d_i . Even better, it is easy to check once again that, with this definition, $d_i d_j = d_{j-1} d_i$ when $i < j$, simply as an evident property of strictly order preserving maps. This is really how we argued for this axiom back in the first place!

So, in summary, the category $\widehat{\Delta}^{op}$ is just the collection of elementary n -simplices together with the face maps (satisfying the face map axiom) and the iterations of face maps. But this should be precisely the prototype for all Delta sets:

Definition 2.9 (Alternative definition for Delta sets). A Delta set is a covariant functor $X : \widehat{\Delta}^{op} \rightarrow \mathbf{Set}$, where $\mathbf{Set}$ is the category of sets and functions. Equivalently, a Delta set is a contravariant functor $\widehat{\Delta} \rightarrow \mathbf{Set}$.

Let's see why this makes sense. A functor takes objects to objects and morphisms to morphisms, and it obeys composition rules. So, unwinding the definition, a covariant functor $\widehat{\Delta}^{op} \rightarrow \mathbf{Set}$ assigns to $[n] \in \widehat{\Delta}^{op}$ a set X_n (which we can think of, and which we refer to, as a set of simplices) and gives us, for each strictly order preserving $[m] \rightarrow [n]$ in $\widehat{\Delta}$ (or its corresponding opposite in $\widehat{\Delta}^{op}$) a generalized face map $X_n \rightarrow X_m$ (which we think of as assigning an m -face to each simplex in X_n). As noted previously, these generalized face maps are all compositions of our standard face maps d_i , so the d_i (and their axioms) are the only ones we usually bother focusing on.

So what just happened? The power of this definition is really in its point of view. Instead of thinking of a Delta set as being made up of a whole bunch of simplices one at a time, we can now think of the standard n -simplex as standing for all of the simplices in X_n , all at once - the functor X assigns to $[n]$ the collection of all of the simplices of X_n (see figure 9). The face map d_i applied to the standard simplex $[n]$ represents all of the i th faces of all the n -simplices simultaneously.




Figure 9: A Delta complex as the functorial image of $\widehat{\Delta}$

At the same, we see how any argument in X really comes from what happens back in $\widehat{\Delta}$. The commutativity axiom $d_i d_j = d_{j-1} d_i$ in a Delta set X is just a consequence of this

being true in the prototype simplex $[n]$ and inherent properties of functors. We'll get a lot of mileage out of this kind of thinking: things we'd like to prove in a Delta set X can often be proven just by proving it in the prototype standard simplex and applying functoriality.

Delta maps. We won't dwell overly long on Delta maps, except to observe that they, too, point toward the need for simplicial sets (however, see [18] where Delta complexes and Delta maps are treated in their own right).

Going directly to the category theoretic definition, given two Delta sets X, Y , thought of as contravariant functors $\hat{\Delta} \rightarrow \mathbf{Set}$, a morphism $X \rightarrow Y$ is a natural transformation of functors from X to Y . Thus a morphism consists of a collection of set maps $X_n \rightarrow Y_n$ that commute with the face maps.

Example 2.10. There is an apparent Delta map from the standard 2-simplex $[0, 1, 2]$ to the cone C of Example 2.6. See Figure 10.




Figure 10: The Delta map from $|\Delta^2|$ to the cone

The astute reader will notice something fishy here. We would hope that simplicial maps of simplicial complexes would yield morphisms of Delta sets. However, consider the collapse $\pi : |\Delta^2| = [0, 1, 2] \rightarrow |\Delta^1| = [0, 1]$ defined by $\pi(0) = 0$ and $\pi(1) = \pi(2) = 1$ (see Figure 5). To be a Delta set morphism, the simplex $[0, 1, 2] \in |\Delta^2|_2$ would have to be taken to an element of $|\Delta^1|_2$. But this set is empty! There are no 2-simplices of $|\Delta^1|$. Something is amiss. We need simplicial sets.

3 Simplicial sets and morphisms

Simplicial sets generalize both simplicial complexes and Delta sets.

When approaching the literature, the reader should be very careful about terminology. Originally ([5]), Delta sets were referred to as *semi-simplicial complexes*, and, once the degeneracy operations we are about to discuss were discovered, the term *complete semi-simplicial complex* (*c.s.s. set*, for short) was introduced. Over time, with Delta sets becoming of less interest, “complete semi-simplicial” was abbreviated back to “semi-simplicial” and eventually to “simplicial,” leaving us with the *simplicial sets* of today. Meanwhile, some modern authors have returned to using “semi-simplicial complexes” to refer to what we are calling Delta sets, on the grounds that, as we will see, the category Δ is the prototype for simplicial sets, not Delta sets, for which we have been using the prototype category $\hat{\Delta}$. This

all sounds very confusing because it is, and the reader is advised to be very careful when reading the literature.²

We try to be careful and use only the three terms “simplicial complex,” “Delta set,” and “simplicial set.” In particular, be sure to note the difference between “simplicial complex” and “simplicial set” going forward.

Degenerate simplices. Recall from Example 2.4 that a simplicial map can collapse a simplex. In that example, we had a simplicial map $\pi : |\Delta^2| \rightarrow |\Delta^1|$ defined on vertices so that $\pi(0) = 0$ and $\pi(1) = \pi(2) = 1$. Recall also that we have begun to think of simplicial complexes and Delta sets as collections of images of standard simplices under appropriate maps. Well, here is a map of the standard 2-simplex $|\Delta^2|$. What image simplex does it give us in $|\Delta^1|$ under π ? In the land of simplicial sets, the image $\pi(|\Delta^2|)$ is an example of a *degenerate* simplex.

Roughly speaking, degenerate simplices are simplices that don’t have the “correct” number of dimensions. A degenerate 3-simplex might be realized geometrically as a 2-dimensional, 1-dimensional, or 0-dimensional object. Geometrically, degenerate simplices are “hidden”; thus the clearest approach to dealing with them lies in the combinatorial notation we have been developing all along.

The key both to the idea and to the notation is in allowing vertices to repeat. The natural way to label $\pi(|\Delta^2|) = \pi([0, 1, 2])$ in our example is as $[0, 1, 1]$, reflecting where the vertices of $|\Delta^2|$ go under the map. This violates our earlier principle that simplices in complexes should be written $[v_0, \dots, v_n]$ with the v_i distinct vertices written in order, but sometimes in mathematics we need a new, more general principle. For degenerate simplices, we’ll keep the orderings but dispense with the uniqueness. Thus, officially, a *degenerate* simplex is a $[v_{i_0}, \dots, v_{i_n}]$ for which the v_{i_j} are *not* all distinct.

Example 3.1. How many 1-simplices, including degenerate ones, are lurking within the elementary 2-simplex $[0, 1, 2]$? A one simplex is still written $[a, b]$, with $a \leq b$, but now repetition is allowed. The answer is six: $[0, 1]$, $[0, 2]$, $[1, 2]$, $[0, 0]$, $[1, 1]$, and $[2, 2]$. See the middle picture in Figure 11.

Similarly, within $|\Delta^2| = [0, 1, 2]$ there are now three kinds of 2-simplices. We have the nondegenerate $[0, 1, 2]$, the 2-simplices that degenerate to 1-dimension such as $[0, 1, 1]$ and $[0, 0, 2]$, and we have the 2-simplices that degenerate to 0-dimensions such as $[0, 0, 0]$ and $[2, 2, 2]$.

Working with degenerate simplices makes drawing diagrams much more difficult. We take a crack at it in Figure 11.

As implied by the diagram, we can think of degenerate simplices as being the images of collapsing maps such as that in Example 2.4.

Of course any simplicial complex or Delta set can be expanded conceptually to include degenerate simplices. In the example of Figure 1, we might have the degenerate 5-simplex $[v_2, v_2, v_2, v_3, v_3]$.

²I thank Jim McClure for explaining to me this historical progression.




Figure 11: The first picture represents all of the 1-simplices in $|\Delta^1|$, including the degenerate ones that are taken to individual vertices. The second picture represents all the 1-simplices in $|\Delta^2|$, and the last picture represents all of the degenerate 2-simplices in $|\Delta^2|$.

Notice also that our innocent little n -dimensional simplicial complexes suddenly contain degenerate simplices of arbitrarily large dimension. Even the 0-simplex $\Delta^0 = [0]$ becomes host to degenerate simplices such as $[0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0]$.

The situation has degenerated indeed! To keep track of it all, we need *degeneracy maps*.

Degeneracy maps. Degeneracy maps are, in some sense, the conceptual converse of face maps. Recall that the face map d_j takes an n -simplex and give us back its j th $(n - 1)$ -face. On the other hand, the j th degeneracy map s_j takes an n -simplex and gives us back the j th degenerate $(n + 1)$ -simplex living inside it.

As usual, we illustrate with the standard n -simplex, which will be a model for what happens in all simplicial sets. Given the standard n -simplex $|\Delta^n| = [0, \dots, n]$, there are $n + 1$ degeneracy maps $s_0, \dots, s_n$, defined by $s_j[0, \dots, n] = [0, \dots, j, j, \dots, n]$. In other words, $s_j[0, \dots, n]$ gives us the unique degenerate $n + 1$ simplex in $|\Delta^n|$ with only the j th vertex repeated.

Again, the geometric concept is that $s_j|\Delta^n|$ can be thought of as the process of collapsing Δ^{n+1} down into Δ^n by the simplicial map π_j defined by $\pi_j(i) = i$ for $i < j$, $\pi_j(j) = \pi_j(j+1) = j$ and $\pi_j(i) = i - 1$ for $i > j + 1$.

This idea extends naturally to simplicial complexes, to Delta sets, and to simplices that are already degenerate. If we have a, possibly degenerate, n -simplex $[v_{i_0}, \dots, v_{i_n}]$ with $i_k \leq i_{k+1}$ for each k , $0 \leq k \leq n$, then we set $s_j[v_{i_0}, \dots, v_{i_n}] = [v_{i_0}, \dots, v_{i_j}, v_{i_j}, \dots, v_{i_n}]$, i.e. repeat

v_{i_j} . This is a degenerate simplex in $[v_{i_0}, \dots, v_{i_n}]$.

It is not hard to see that any degenerate simplex can be obtained from an ordinary simplex by repeated application of degeneracy maps. Thus, just as any face of a simplex can be obtained by using compositions of the d_i , any degenerate simplex can be obtained from compositions of the s_i .

Also, as for the d_i , there are certain natural relations that the degeneracy maps possess. In particular, if $i \leq j$, then $s_i s_j[0, \dots, n] = [0, \dots, i, i, \dots, j, j, \dots, n] = s_{j+1} s_i[0, \dots, n]$. Note that we have $j + 1$ in the last formula, not s_j , since the application of s_i pushes j one slot to the right.

Furthermore, there are relations amongst the face and degeneracy operators. These are a little more awkward to write down since there are three possibilities:

$$\begin{aligned} d_i s_j &= s_{j-1} d_i & \text{if } i < j, \\ d_j s_j &= d_{j+1} s_j = \text{id}, \\ d_i s_j &= s_j d_{i-1} & \text{if } i > j + 1. \end{aligned}$$

These can all be seen rather directly. For example, applying either side of the first formula to $[0, \dots, n]$ yields $[0, \dots, \hat{i}, \dots, j, j, \dots, n]$. Note also that the middle formula takes care of both $i = j$ and $i = j + 1$.

Simplicial sets. We are finally ready for the definition of simplicial sets:

Definition 3.2. A *simplicial set* consists of a sequence of sets $X_0, X_1, \dots$ and, for each $n \geq 0$, functions $d_i : X_{n+1} \rightarrow X_n$ and $s_i : X_n \rightarrow X_{n+1}$ for each i with $0 \leq i \leq n + 1$ such that

$$\begin{aligned} d_i d_j &= d_{j-1} d_i & \text{if } i < j, \\ d_i s_j &= s_{j-1} d_i & \text{if } i < j, \\ d_j s_j &= d_{j+1} s_j = \text{id}, \\ d_i s_j &= s_j d_{i-1} & \text{if } i > j + 1, \\ s_i s_j &= s_{j+1} s_i & \text{if } i \leq j. \end{aligned} \tag{1}$$

Example 3.3. Every simplicial complex or Delta set can be made into a simplicial set by adjoining all possible degenerate simplices. Conversely, each simplicial set is also a Delta set by neglect of structure. However, a simplicial set does not necessarily come from a simplicial complex as, for example, not every Delta set is a simplicial complex.

Example 3.4. The standard 0-simplex $X = [0]$, now thought of as a simplicial set, is the unique simplicial set with one element in each X_n , $n \geq 0$. The element in dimension n is $\overbrace{[0, \dots, 0]}^{n+1 \text{ times}}$.

Example 3.5. As a simplicial set, the standard 1-simplex $X = [0, 1]$ already has $n + 2$ elements in each X_n . For example, $X_2 = \{[0, 0, 0], [0, 0, 1], [0, 1, 1], [1, 1, 1]\}$.

Remark 3.6. We will use Δ^n or $[0, \dots, n]$ to refer to the standard n -simplex, thought of as a simplicial set.

Example 3.7. Now for an example familiar from algebraic topology. Given a topological space X , let $\mathcal{S}(X)_n$ be the set of continuous functions from $|\Delta^n|$ to X . Together with face and degeneracy maps that we will describe in a moment, these constitute a simplicial set called the singular set of X . The singular chain complex $S_*(X)$ from algebraic topology has each $S_n(X)$ equal to the free abelian group generated by $\mathcal{S}(X)_n$.

To define the face and degeneracy maps, let $\sigma : |\Delta^n| \rightarrow X$ be a continuous map representing a singular simplex (Figure 12). The singular simplex $d_i\sigma$ is defined as the restriction of σ to the i th face of $|\Delta^n|$. Equivalently it is the composition of σ and the simplicial inclusion map $[0, \dots, n-1] \rightarrow [0, \dots, \hat{i}, \dots, n]$ (Figure 13). These are precisely the same as the terms that show up in the boundary map of the singular chain complex where $\partial = \sum_{i=0}^n (-1)^i d_i$.




Figure 12: A singular simplex




Figure 13: A face of a singular simplex

On the other hand, the degeneracy s_i takes the singular simplex σ to the composition of $\sigma : |\Delta^n| = [0, \dots, n] \rightarrow X$ with the geometric collapse represented by the degeneracy

$[0, \dots, n+1] \rightarrow [0, \dots, i, i, \dots, n]$. Once again, a degenerate simplex is a collapsed version of an ordinary simplex (Figure 14).




Figure 14: A degenerate singular simplex

$\mathcal{S}(X)$ is clearly a simplicial set: the face maps take a singular simplex to the map defined along the face of the model simplex and the degeneracy is represented by a collapsed simplex. It is our usual model, just redesigned within the context of the continuous map σ .

Some more example of simplicial sets are given below in Section 4, where we can better study their geometric manifestations.

Nondegenerate simplices.

Definition 3.8. A simplex $x \in X_n$ is called *nondegenerate* if x cannot be written as $s_i y$ for any $y \in X_{n-1}$ and any i .

Every simplex in the sense of Section 2 of a simplicial complex or Delta set is a nondegenerate simplex of the corresponding simplicial set. If Y is a topological space, an n -simplex of $\mathcal{S}(Y)$ is nondegenerate if it cannot be written as the composition $\Delta^n \xrightarrow{\pi} \Delta^k \xrightarrow{\sigma} Y$, where π is a simplicial collapse with $k < n$ and σ is a singular k -simplex.

Note that it is possible for a nondegenerate simplex to have a degenerate face (see Example 4.7, below, though it might be good practice to try to come up with your own example first). It is also possible for a degenerate simplex to have a nondegenerate face (for example, we know $d_j s_j x = x$ for any x , degenerate or not).

The categorical definition. As for Delta sets, the basic properties of simplicial sets derive from those of the standard n -simplex. In fact, that is where the prototypes of both the face and degeneracy maps live and where we first developed the axioms relating them. Thus it is not surprising (at this point) that there is a categorical definition of simplicial sets, analogous to the one for Delta sets, in which each simplicial set is the functorial image of a category, Δ , built from the standard simplices.

Definition 3.9. The category Δ has as objects the finite ordered sets $[n] = \{0, 1, 2, \dots, n\}$. The morphisms of Δ are order preserving functions $[m] \rightarrow [n]$.

Notice that the only difference between the definitions of $\widehat{\Delta}$ and Δ is that the morphisms in Δ only need to be order preserving and not strictly order preserving. Thus, equating the objects $[n]$ with the simplices Δ^n , the morphisms no longer need to represent only inclusions of simplices but may represent degeneracies as well. In more familiar notation, a typical morphism, say, $f : [5] \rightarrow [3]$ might be described by $f[0, 1, 2, 3, 4, 5] = [0, 0, 2, 2, 2, 3]$, which can be thought of as the simplicial complex map taking the 5-simplex degenerately to the 2-face of the 3-simplex spanned by 0, 2, and 3.

As in $\widehat{\Delta}$, the morphisms in Δ are generated by certain maps between neighboring cardinalities $D_i : [n] \rightarrow [n+1]$ and $S_i : [n+1] \rightarrow [n]$, $0 \leq i \leq n$. The D_i are just as for $\widehat{\Delta}$: $D_i[0, \dots, n] = [0, \dots, i, \dots, n+1]$. The new maps, which couldn't exist in $\widehat{\Delta}$, are defined by $S_i[0, \dots, n+1] = [0, \dots, i, i, \dots, n]$. It is an easy exercise to verify that all morphisms in Δ are compositions of the D_i and S_i and that these satisfy axioms analogous to those in the definition of simplicial set. Later on, we will also use D_i and S_i to stand for the geometric maps they induce on the standard geometric simplices.

To get to our categorical definition of simplicial sets, we must, as for Delta sets, consider Δ^{op} . The maps D_i become their opposites, denoted d_i , and these correspond to the face maps as before: the opposite of the inclusion $D_i : [n] \rightarrow [n+1]$ of the i th face is the i th face map, d_i , which assigns to the n -simplex its i th face. The opposites of the S_i become the degeneracies; the opposite of the collapse $S_i : [n+1] \rightarrow [n]$ that pinches together the i and $i+1$ vertices of an $n+1$ simplex is the i th degeneracy map, s_i , which assigns to the n -simplex Δ^n the degenerate $n+1$ -simplex within Δ^n that repeats the i th vertex. See figure 15.




Figure 15: How to visualize D_i , d_i , S_i , and s_i . Our difficulty with drawing degeneracies extends here so that we represent the image of s_i pictorially by the picture for S_i . In other words, the image of s_1 in the bottom right is the degenerate 2-simplex arising from the collapse map S_1 .

Of course, the d_i and s_i can be checked to satisfy the axioms in the definition of simplicial set given above.

Definition 3.10 (Categorical definition of simplicial set). A *simplicial set* is a contravariant functor $X : \Delta \rightarrow \mathbf{Set}$ (equivalently, a covariant functor $X : \Delta^{op} \rightarrow \mathbf{Set}$).

The reader should compare this with the categorical definition of Delta sets and reassure himself/herself that this definition is equivalent to Definition 3.2. As for Delta sets, the power in this definition is that we can think of the standard n -simplex as standing for all of the simplices in X_n , all at once - the functor X assigns to $[n]$ all of the n -simplices in X_n - and the standard face and degeneracy maps d_i and s_i of $[n] = \Delta^n$ pick out all of the faces and degeneracies of X_n by functoriality.

Example 3.11. Let's re-examine the singular set $\mathcal{S}(Y)$ of the topological space Y from this point of view. The singular set $\mathcal{S}(Y)$ is a functor $\Delta^n \rightarrow \mathbf{Set}$ that assigns to $[n]$ the set $\mathrm{Hom}_{\mathbf{Top}}(\Delta^n, Y)$, the set of all continuous maps from Δ^n to Y . The face and degeneracy maps are completely determined by the face and degeneracy maps of Δ and the properties of functors, which require that the following diagrams commute:

$$\begin{array}{ccc} [n] & \longrightarrow & \mathrm{Hom}_{\mathbf{Top}}(\Delta^n, Y) \\ d_i \downarrow & & \downarrow d_i \\ [n-1] & \longrightarrow & \mathrm{Hom}_{\mathbf{Top}}(\Delta^{n-1}, Y) \end{array} \qquad \begin{array}{ccc} [n] & \longrightarrow & \mathrm{Hom}_{\mathbf{Top}}(\Delta^n, Y) \\ s_i \downarrow & & \downarrow s_i \\ [n+1] & \longrightarrow & \mathrm{Hom}_{\mathbf{Top}}(\Delta^{n+1}, Y). \end{array}$$

The reader should check that the definitions for the face and degeneracy maps of the singular set defined above indeed lead to commutativity of the diagram. (Notice that the maps on the right sides of these squares should more appropriately be labeled $\mathcal{S}(Y)(d_i)$ and $\mathcal{S}(Y)(s_i)$, but we stick with common practice and use d_i and s_i for face and degeneracy maps wherever we find them.)

Simplicial morphisms. Simplicial sets themselves constitute a category $\mathbf{S}$. In the categorical language, if X and Y are simplicial sets (and thus functors $\Delta \rightarrow \mathbf{Set}$), then a morphism $f : X \rightarrow Y$ is a natural transformation of functors. Unwinding this to more concrete language, f consists of set maps $f_n : X_n \rightarrow Y_n$ that commute with face operators and with degeneracy operators.

Example 3.12. At last we have a context in which to explain properly the collapse map $\pi : |\Delta^2| \rightarrow |\Delta^1|$ of Example 2.4. We extend π to the map of simplicial sets $\pi : \Delta^2 \rightarrow \Delta^1$ that takes $\Delta^2 = [0, 1, 2]$ to the degenerate simplex $[0, 1, 1] = s_1([0, 1])$. At the same time, it is doing an infinite number of other things. For example, it takes the vertex $[0] \in \Delta^2$ to $[0] \in \Delta^1$, it takes the vertices $[1], [2] \in \Delta^2$ to $[1] \in \Delta^1$, it takes the 1-simplex $[0, 1] \in \Delta^2$ to $[0, 1] \in \Delta^1$, it takes the 1-simplex $[1, 2] \in \Delta^2$ to the degenerate 1-simplex³ $[1, 1] = s_0[1] \in \Delta^1$, and it even takes the degenerate simplex $[0, 1, 1, 2, 2, 2] = s_4 s_3 s_1[0, 1, 2] \in \Delta^2$ to the degenerate simplex $s_4 s_3 s_1[0, 1, 1] = [0, 1, 1, 1, 1, 1] \in \Delta^1$. And much much more.

³Careful: $[1]$ is a 0-simplex, so s_0 is the appropriate (indeed the only well-defined) degeneracy map. Remember that s_0 tells us to repeat what occurs in the 0th place - it doesn't know what's in that place.

Example 3.13. Notice that, unlike simplicial maps on simplicial complexes, simplicial morphisms on simplicial sets are not completely determined by what happens on vertices. For example, consider the simplicial morphisms from Δ^1 to the Delta set decomposition of the circle with two unique vertices of Example 2.7. If we have a simplicial morphism that takes $[0]$ to $[v_0]$ and $[1]$ to $[v_1]$, there are still two possibilities for where to send $[0, 1]$.

Example 3.14. On the other hand, given a map of *ordered* simplicial complexes $f : X \rightarrow Y$, this induces a map of the associated simplicial sets obtained by adjoining all naturally occurring degenerate simplices, ie. all simplices of the form $[v_{i_0}, \dots, v_{i_n}]$ with $i_0 \leq \dots \leq i_n$, where all v_{i_j} are vertices of some particular simplex of X (or, respectively, of Y). In this case, a function on vertices *does* determine a simplicial map.

Remark 3.15. Notice that it is always enough to define a simplicial morphism by what it does to nondegenerate simplices. What happens to the degenerate simplices is forced by the definition since, e.g. $f(s_i(x)) = s_i(f(x))$. Similarly, what happens on faces is forced by what happens on the simplices of which they are faces. Thus, altogether, simplicial morphisms can be described by specifying what they do to a rather small collection of nondegenerate simplices.

From here on, we'll abandon the distinction between “simplicial map” and “simplicial morphism” and use the terms interchangeably as applied to simplicial sets.

4 Realization

If the idea of simplicial objects is to abstract from geometry to combinatorics, there should be a way to reverse that process and turn simplicial sets into geometric objects. Indeed that is the case. The definition looks a bit off-putting at first (what concerning simplicial sets doesn't?), but, in fact, we'll see that simplicial realization is a very natural thing to do.

Definition 4.1. Let X be a simplicial set. Give each set X_n the discrete topology and let Δ^n be the n -simplex with its standard topology. The *realization* $|X|$ is given by

$$|X| = \coprod_{n=0}^{\infty} X_n \times |\Delta^n| / \sim,$$

where $\sim$ is the equivalence relation generated by the relations $(x, D_i(p)) \sim (d_i(x), p)$ for $x \in X_{n+1}, p \in \Delta^n$ and the relations $(x, S_i(p)) \sim (s_i(x), p)$ for $x \in X_{n-1}, p \in \Delta^n$. Here D_i and S_i are the face inclusions and collapses induced on the standard geometric simplices as in our discussion above of the category Δ .

To see why this definition makes sense, let's think about how we would like to form a simplicial complex out of the data of a simplicial set. From the get-go, we have been thinking of the X_n as collections of simplices. So this is just what $X_n \times |\Delta^n|$ is: a disjoint collection of simplices, one for each element of X_n . The next natural thing to do is to identify common faces. This is precisely what the relation $(x, D_i(p)) \sim (d_i(x), p)$ encodes (see Figure 16): The first term of $(x, D_i(p)) \subset (x, |\Delta^{n+1}|)$ is an $n+1$ simplex of X and the second term $D_i(p)$ is a point on the i th face of a geometric $n+1$ simplex. On the other hand, $(d_i(x), p)$ is the i th

face of x together with the same point, now in a stand-alone n -simplex. So the identification described just takes the n -simplex corresponding to $d_i(x)$ in $X_n \times |\Delta^n|$ and glues it as the i th face of the $n + 1$ simplex assigned to x in $X_{n+1} \times |\Delta^{n+1}|$. Since a similar gluing is done for any other y and j such that $d_j(y) = d_i(x)$, the effect is to glue faces of simplices together.




Figure 16: In the realization, the 1-simplex representing $d_0 x$, pictured on the right, is glued to the 2-simplex representing x , pictured on the left, along the appropriate face

The next natural thing to do is suppress the degenerate simplices, since they're encoded within nondegenerate simplices anyway. This is what the relation $(x, S_i(p)) \sim (s_i(x), p)$ for $x \in X_{n-1}, p \in |\Delta^n|$ does, although more elegantly. This relation tells us that given a degenerate n -simplex $s_i(x)$ and a point p in the pre-collapse n -simplex $|\Delta^n|$, we should glue p to the $n - 1$ simplex represented by x at the point $S_i(p)$ in the image of the collapse map. That still sounds a little confusing, but the idea is straightforward: $|\Delta^n|$ s corresponding to degenerate n -simplices get collapsed in the natural way into the $n - 1$ -simplices they are degeneracies of. See Figure 17. We note also that there is no reason to believe that x itself is nondegenerate. It might be, in which case the simplex corresponding to x is itself collapsed. This provides no difficulty.




Figure 17: In the realization, the 2-simplex representing $s_1 x$, pictured on the right, is glued to the 1-simplex representing x , pictured on the left, via the appropriate collapse, depicted by S_1

Example 4.2. Recall that the 0-simplex $[0]$, thought of as a simplicial set, has one simplex $[0, \dots, 0]$ in each dimension ≥ 0 . Thus $||[0]|| = \coprod_{i=0}^{\infty} |\Delta^i| / \sim$. So in dimension 0 we have a single vertex v . In dimension 1, we have a single simplex $[0, 0] = s_0[0]$. The gluing instructions tell us to identify each $(s_i(x), p) = ([0, 0], p) \in ([0, 0], |\Delta^1|)$ with $([0], S_0(p)) = ([0], v)$. Thus the $|\Delta^1|$ in dimension 1 gets collapsed to the vertex. Similarly, since each point of the 2-simplex

$([0, 0, 0], |\Delta^2|)$ gets identified to a point of $([0, 0], |\Delta^1|)$, and so on, we see that the whole situation collapses down to a single vertex.

Example 4.3. Generalizing the preceding example, $|[0, \dots, n]| = |\Delta^n|$ is just the standard geometric n -simplex, justifying our earlier use of notation. We encourage the reader to explore this example on his or her own, noting that all of the degenerate simplices wind up tucked away within actual faces of $|\Delta^n|$, just where we expect them.

Example 4.4. More generally, given any simplicial complex, the realization of the simplicial set associated to it by adjoining only degenerate simplices returns the original simplicial complex.

Example 4.5. There is an analogous realization procedure for Delta sets. Given a Delta set X , we can define the realization $|X|$ by

$$|X| = \prod_{n=0}^{\infty} X_n \times |\Delta^n| / \sim,$$

where $\sim$ is the equivalence relation generated by $(x, D_i(p)) \sim (d_i(x), p)$ for $x \in X_{n+1}, p \in \Delta^n$. These realizations yield the types of spaces we have been drawing already to represent Delta sets. These are sometimes called Delta complexes; see, e.g., [9].

However, given a simplicial set X , the simplicial set realization of X is not generally going to be the same as the Delta set realization of the associated Delta set, say X_{Δ} , that we obtain by neglect of structure.

For example, consider the simplicial set Δ^0 . As noted, its simplicial realization, $|\Delta^0|$ is the topological space consisting of a single point. But recall that the simplicial set Δ^0 has exactly one simplex in each dimension, and the neglect of structure that turns this into a Delta set drops the degeneracy relation but still leaves a Delta set with one simplex in each dimension and all face maps the unique possible ones. Thus the Delta set realization $|\Delta^0|_{\Delta}$ is an infinite dimensional CW complex with one cell in each dimension whose n -dimensional cell is attached by gluing each face of an n -simplex, in an order preserving manner, to the image of the unique $n - 1$ simplex in the $n - 1$ skeleton. Thus the 1-skeleton of $|\Delta^0|_{\Delta}$ is a circle, the 2-skeleton is the “dunce cap” (see, e.g., [2, Section 14]), and so on. This is evidently not homeomorphic to Δ^0 . (However, they are homotopy equivalent; this will be true in general, see [18]).

In what follows, discussion of “realization” and the notation $|X|$ will refer exclusively to simplicial set realization unless noted otherwise.

Example 4.6. Let Y be a topological space, and let $\mathcal{S}(Y)$ be its singular set. $|\mathcal{S}(Y)|$ will be huge (unless Y is discrete - what will it be then?). While this looks discouraging, it turns out that $|\mathcal{S}(Y)|$ is homotopy equivalent to Y . Thus, for the purposes of algebraic topology, Y and $|\mathcal{S}(Y)|$ are indistinguishable. So perhaps, wearing the appropriate glasses, Y and $\mathcal{S}(Y)$ can be thought of as the same thing? We’ll get back to this idea.

Example 4.7. As noted in Example 4.4, the realization of a simplicial set that we obtained from a simplicial complex is the original simplicial complex. So, for example, we can obtain a topological $n - 1$ sphere as the realization of the boundary of the n -simplex, $\partial\Delta^n$. What does $\partial\Delta^n$ look like as a simplicial set? Since every m -simplex of $\partial\Delta^n$ is also a simplex of

Δ^n , each can be written $[i_0, \dots, i_m]$, where $0 \leq i_0 \leq \dots \leq i_m \leq n$. The only caveat is that we cannot allow any m -simplex that contains all of the vertices $0, \dots, n$, since any such simplex would either be the “top face” $[0, \dots, n]$, itself, or a degeneration of it, and these are not allowable faces of $\partial\Delta^n$. In summary, then, we have $S^{n-1} \cong |\partial\Delta^n|$, where $\partial\Delta^n$ is the simplicial set consisting of all sequences of the numbers $0, \dots, n$ that do not contain all of the numbers $0, \dots, n$.

Is this the most efficient way to realize S^{n-1} as the realization of a simplicial set? After all, $\partial\Delta^n$ contains quite a number of simplices, many of which are nondegenerate (the interested reader might go and count them). Here is another way to do it, at least for $n \geq 2$, suggested by CW complexes. Let X be a simplicial set whose only nondegenerate simplices are denoted by $[0] \in X_0$ and $[0, \dots, n-1] \in X_{n-1}$. All simplices in X_i , $0 < i < n-1$, are the degenerate simplices $[0, \dots, 0]$. This, of course, forces all of the faces of $[0, \dots, n-1]$ to be $[0, \dots, 0]$, and we see that the realization $|X|$ is equivalent to the standard construction of S^{n-1} as a CW complex by collapsing the boundary of an $n-1$ cell to a point. See Figure 18.




Figure 18: The realization of the simplicial set consisting of only two nondegenerate simplices, one in dimension 0 and the other in dimension 2, is the sphere S^2 ; this picture represents the image of the nondegenerate simplex of dimension 2 in the realization

The preceding example is instructive on several different points:

1. The second part of Example 4.7 relies strongly on the existence of degenerate simplices. For $n > 2$, we cannot construct S^{n-1} this way as the realization of a Delta set. A Delta set with an $n-1$ simplex would require actual (nondegenerate) $n-2$ simplices as its faces. Of course we can still get S^{n-1} as the realization of the Delta set corresponding to $\partial\Delta^n$.
2. Notice that the realization of a simplicial set does not necessarily inherit the structure of a simplicial complex, at least not in any obvious way from the data of the simplicial set.
3. Realizations are non-unique, in the sense that very different looking simplicial sets can have the same geometric realization up to homeomorphism. This is not surprising, since there are many ways to triangulate a piecewise-linear space.

Example 4.7 is also disconcerting in that the reader may be getting worried that realizations of simplicial sets might be very complicated to understand with all of the gluing and collapsing that can occur. To mitigate these concerns somewhat, we first observe that all degenerate simplices do get collapsed down into the simplices of which they are degeneracies, and so constructing a realization depends only on understanding what happens to the nondegenerate simplices. A second concern would be that two nondegenerate simplices might be glued together. This would happen if it were possible for two nondegenerate simplices to have a common degeneracy. Luckily, this does not happen, as we demonstrate in the following proposition. As a corollary, we can conclude that the realization of a simplicial simplex is made up of the disjoint unions of the interiors of the nondegenerate simplices. We must limit this statement to the interiors as the faces of a nondegenerate simplex may be degenerate, as in the second part of Example 4.7 - meanwhile, nondegenerate faces will look out for themselves!

Proposition 4.8. *A degenerate simplex is a degeneracy of a unique nondegenerate simplex. In other words, if z is a degenerate simplex, then there is a unique nondegenerate simplex x such that $z = s_{i_0} \cdots s_{i_k} x$, for some collection of degeneracy maps $s_{i_0}, \dots, s_{i_k}$.*

Proof. First note that if we have a z such that $z = f_{i_0} \cdots f_{i_k} y$ for some nondegenerate y , where each f_{i_j} is either a face or degeneracy map, then we could trade the face maps to the right using the simplicial set axioms and then replace y with one of its proper faces. If the face happens to be degenerate, we can repeat the process. Eventually this procedure halts. So a degenerate simplex can always be written in the form $s_{i_0} \cdots s_{i_k} x$.

Suppose now x and y are nondegenerate simplices, possibly of different dimensions, and that $Sx = Ty$, where S and T are compositions of degeneracy operators. Now, suppose $S = s_{i_0} \cdots s_{i_k}$. Let $D = d_{i_k} \cdots d_{i_0}$. Then $x = DSx = DTy$, using the simplicial set axioms for the first equality, and, again by trading to the right, we obtain $x = \tilde{T}\tilde{D}y$ for some composition of face operators $\tilde{D}$ and some composition of degeneracies $\tilde{T}$. But, by hypothesis, x is nondegenerate, so $\tilde{T}$ must be vacuous, and we must have $x = \tilde{D}y$. That is x is a face of y . But we could repeat the argument reversing x and y to obtain that y is also face of x . But this is impossible unless $x = y$. $\square$

Another comforting fact is the following theorem:

Theorem 4.9. *If X is a simplicial set, then $|X|$ is a CW complex with one n -cell for each nondegenerate n -simplex of X .*

Proof. The first part of the proof is the preceding proposition. For the remainder of the proof, we refer to Milnor's notes on realization, which can be found in [13] (or, alternatively, to [11, Theorem 14.1]). $\square$

The adjointness relation. The realization functor $|\cdot|$ turns out to be adjoint to the singular simplex functor $\mathcal{S}(\cdot)$.

Theorem 4.10. *If X is a simplicial set and Y is a topological space, then*

$$\text{Hom}_{\text{Top}}(|X|, Y) \cong \text{Hom}_{\mathbf{S}}(X, \mathcal{S}(Y)).$$

Sketch of proof. We identify the maps $\Psi : \text{Hom}_{\mathbf{Top}}(|X|, Y) \rightarrow \text{Hom}_{\mathbf{S}}(X, \mathcal{S}(Y))$ and $\Phi : \text{Hom}_{\mathbf{S}}(X, \mathcal{S}(Y)) \rightarrow \text{Hom}_{\mathbf{Top}}(|X|, Y)$ and leave it to the reader both to check carefully that these are well-defined and to show that they are mutual inverses.

A map $f \in \text{Hom}_{\mathbf{S}}(X, \mathcal{S}(Y))$ assigns to each n -simplex $x \in X$ a continuous function $\sigma_x : |\Delta^n| \rightarrow Y$. Let $\Phi(f)$ be the continuous function that acts on the simplex $(x, |\Delta^n|) \in |X|$ by applying σ_x to $|\Delta^n|$.

Conversely, given a function $g \in \text{Hom}_{\mathbf{Top}}(|X|, Y)$, then the restriction of g to a nondegenerate simplex $(x, |\Delta^n|)$ yields a continuous function $|\Delta^n| \rightarrow Y$ and thus an element of $\mathcal{S}(Y)_n$. If $(x, |\Delta^n|)$ represents a degenerate simplex, then we precompose with the appropriate collapse map of Δ^n into $|X|$ before applying g . $\square$

One can say much more on the relation between simplicial sets and topology categories. For example, see Theorem 10.1 below, according to which the homotopy category of CW complexes is equivalent to the homotopy category of simplicial sets satisfying a condition called the *Kan condition*. The Kan condition is defined in Section 7.

5 Products

Before we move on to a look at simplicial homotopy, we will need to know about products of simplicial sets. For those accustomed to products of simplicial complexes or products of chain complexes, the definition of the product of simplicial sets looks surprisingly benign by comparison.

Definition 5.1. Let X and Y be simplicial sets. Then their product $X \times Y$ is defined by

1. $(X \times Y)_n = X_n \times Y_n = \{(x, y) \mid x \in X_n, y \in Y_n\}$,
2. if $(x, y) \in (X \times Y)_n$, then $d_i(x, y) = (d_i x, d_i y)$,
3. if $(x, y) \in (X \times Y)_n$, then $s_i(x, y) = (s_i x, s_i y)$.

Notice that there are evident projection maps $\pi_1 : X \times Y \rightarrow X$ and $\pi_2 : X \times Y \rightarrow Y$ given by $\pi_1(x, y) = x$ and $\pi_2(x, y) = y$. These maps are clearly simplicial morphisms.

Definition 5.1 looks disturbingly simple-minded, but it is vindicated by the following important theorem.

Theorem 5.2. *If X and Y are simplicial sets, then $|X \times Y| \cong |X| \times |Y|$ (in the category of compactly generated Hausdorff spaces). In particular, if X and Y are countable or if one of $|X|, |Y|$ is locally finite as a CW complex, then $|X \times Y| \cong |X| \times |Y|$ as topological spaces.*

We refer the reader to [11, Theorem 14.3] or [12] for a proof in the latter situations and to [7, Chapter III] for a proof of the general case. However, since an example is perhaps worth a thousand proofs, we have a detailed look at some special cases.

Example 5.3. Let X be any simplicial set, and let $Y = \Delta^0 = [0]$. Since Δ^0 has a unique element in each dimension, $X \times \Delta^0 \cong X$. So indeed, $|X \times \Delta^0| \cong |X| \times |\Delta^0| \cong |X|$.

Example 5.4. The first interesting example is $\Delta^1 \times \Delta^1$. We would like to see that $|\Delta^1 \times \Delta^1| \cong |\Delta^1| \times |\Delta^1|$, the square. As discussed in Section 4, we need to focus on the nondegenerate simplices of $\Delta^1 \times \Delta^1$. The reader can refer to Figure 19 for the following discussion.




Figure 19: The realization of $\Delta^1 \times \Delta^1$

First, in dimension 0, we have the product 0-simplices

$$X_0 = \{([0], [0]), ([1], [0]), ([0], [1]), ([1], [1])\},$$

the four vertices of the square.

In dimension 1, we have the pairs (e, f) , where e and f are 1-simplices of Δ^1 . There are three possibilities for each of e and f - $[0, 0]$, $[0, 1]$, and $[1, 1]$. So there are nine 1-simplices of $\Delta^1 \times \Delta^1$.

There is only one 1-simplex that is made up completely of nondegenerate simplices: $([0, 1], [0, 1])$. Since $d_0([0, 1], [0, 1]) = (0, 0)$ and $d_1([0, 1], [0, 1]) = (1, 1)$, the simplex $([0, 1], [0, 1])$ must be the diagonal. Those with one nondegenerate and one degenerate 1-simplex are $([0, 0], [0, 1])$, $([0, 1], [0, 0])$, $([1, 1], [0, 1])$ and $([0, 1], [1, 1])$, which, as we see by checking the endpoints, are respectively the left, bottom, right, and top of the square. The other four 1-simplices are the degeneracies of the vertices. For example, $([0, 0], [1, 1]) = (s_0[0], s_0[1]) = s_0([0], [1])$.

Now for the 2-simplices - here's where things get a little tricky. There are four 2-simplices of Δ^1 : $[0, 0, 0]$, $[0, 0, 1]$, $[0, 1, 1]$, and $[1, 1, 1]$. So there are sixteen 2-simplices of $\Delta^1 \times \Delta^1$. There are two possible degeneracy maps, s_0 and s_1 , from $(\Delta^1 \times \Delta^1)_1$ to $(\Delta^1 \times \Delta^1)_2$. These act on the nine 1-simplices, but there are not eighteen degenerate 2-simplices since $s_0 s_0 = s_1 s_0$, and we know there are four degenerate 1-simplices $s_0 v_i$ of $\Delta^1 \times \Delta^1$ corresponding to the degeneracies of the four vertices. Removing these redundancies leaves fourteen degenerate simplices. There are no other redundancies since $s_0 s_0 = s_1 s_0$ is the only relation on s_1 and s_0 . The remaining two 2-simplices are nondegenerate. These turn out to be $([0, 0, 1], [0, 1, 1])$ and $([0, 1, 1], [0, 0, 1])$, which are the two triangles, as one can check by computing face maps.

Next, we need to see that all 3-simplices of $\Delta^1 \times \Delta^1$ and above are degenerate. We first observe that each 3-simplex of Δ^1 must be a double degeneracy of a 1-simplex (since there are no nondegenerate simplices of Δ^1 of dimension greater than 1). But there are

only six such options, of the forms s_0s_0e , s_0s_1e , s_1s_0e , s_1s_1e , s_2s_0e and s_2s_1e for a (possibly degenerate) 1-simplex e . However, the simplicial set axioms reduce this to the possibilities s_1s_0e , s_2s_0e , and s_2s_1e . But then, again by the axioms,

$$\begin{aligned}
(s_1s_0e, s_1s_0f) &= s_1(s_0e, s_0f) \\
(s_1s_0e, s_2s_0f) &= (s_0s_0e, s_0s_1f) = s_0(s_0e, s_1f) \\
(s_1s_0e, s_2s_1f) &= (s_1s_0e, s_1s_1f) = s_1(s_0e, s_1f) \\
(s_2s_0e, s_1s_0f) &= (s_0s_1e, s_0s_0f) = s_0(s_1e, s_0f) \\
(s_2s_0e, s_2s_0f) &= s_2(s_0e, s_0f) \\
(s_2s_0e, s_2s_1f) &= s_2(s_0e, s_1f) \\
(s_2s_1e, s_1s_0f) &= (s_1s_1e, s_1s_0f) = s_1(s_1e, s_0f) \\
(s_2s_1e, s_2s_0f) &= s_2(s_1e, s_0f) \\
(s_2s_1e, s_2s_1f) &= s_2(s_1e, s_1f).
\end{aligned}$$

So all 3-simplices of $\Delta^1 \times \Delta^1$ are degenerate. It also follows that all higher dimension simplices are degenerate: the terms in any such product must be further degeneracies of these particular doubly degenerate 1-simplices, and using the simplicial set axioms, we can move s_0 and s_1 to the left in all expressions. Then we can proceed as in the above list of computations.

That last bit isn't very intuitive, but the low-dimensional part makes some sense. If we take the product of two CW complexes, the cells of the product will be the products of the cells $C_1 \times C_2$, where C_1 and C_2 are not necessarily of the same dimension. These mixed dimensional cells occur here as products of nondegenerate simplices with degenerate simplices. What makes matters difficult is that we must preserve a simplicial structure. This forced "triangulation" is what makes matters somewhat complicated.

It will also be useful for us to look more closely at the products $\Delta^p \times \Delta^q$. After all, all products will be made up of these building blocks. The main point of interest for us is that the simplicial product construction yields the same triangulation structure that may be familiar from homotopy arguments in courses in beginning algebraic topology.

Example 5.5. Since we know that $|\Delta^p \times \Delta^q| = |\Delta^p| \times |\Delta^q|$, let us focus on the nondegenerate $p + q$ simplices of $\Delta^p \times \Delta^q$. We let E_j stand for the unique non-degenerate j -simplex of Δ^j . We note immediately that any nondegenerate $p + q$ simplex s of $\Delta^p \times \Delta^q$ (and hence the only ones that appear nondegenerately in the realization) must have the form $s = (SE_p, S'E_q)$, where S and S' are sequences of degeneracy maps. Why? Otherwise s would have to be of the form $s = (\bar{S}t, \bar{S}'t')$, where $\bar{S}$ and $\bar{S}'$ are again sequences of degeneracy maps and t and t' are faces of E_p and E_q , respectively. But in this case, we would have $s \in F \times F'$, where F and F' are faces of Δ^p and Δ^q . Consequently the image of $s \times |\Delta^{p+q}|$ in the realization of $\Delta^p \times \Delta^q$ will in fact lie within the realization $|F| \times |F'|$. But then $\dim |s| < p + q$, so s could not have been a nondegenerate $p + q$ simplex.

So now we see that $s = (SE_p, S'E_q)$, and for dimensional reasons, we can write this as $s = (s_{i_q} \dots s_{i_1} E_p, s_{j_p} \dots s_{j_1} E_q)$. Furthermore, using the simplicial set axioms, we can assume

that $0 \leq i_1 < \dots < i_q < p + q$ and $0 \leq j_1 < \dots < j_q < p + q$. Now notice that the collection $\{i_1, \dots, i_q, j_1, \dots, j_q\}$ consists of $p + q$ numbers from 0 to $p + q - 1$. Furthermore, there can be no redundancy, since if $i_k = j_{k'}$ for some k and k' , then again by the axioms, we can pull these indices to the front to get $s = (s_{i_k} \tilde{S} E_p, s_{j_{k'}} \tilde{S}' E_q) = s_{i_k} (\tilde{S} E_p, \tilde{S}' E_q)$ for some $\tilde{S}, \tilde{S}'$, making s degenerate.

Thus we conclude that the nondegenerate $p + q$ simplices of $\Delta^p \times \Delta^q$ are precisely those of the form $s = (s_{i_q} \dots s_{i_1} E_p, s_{j_p} \dots s_{j_1} E_q)$, where the i_k and j_k are increasing series of integers from 0 to $p + q - 1$, all completely distinct.

In the special case $\Delta^p \times \Delta^1 = \Delta^p \times I$, this rule for nondegenerate $p + 1$ dimensional simplices reduces to the form $s = (s_i E_p, s_{j_p} \dots s_{j_1} e)$, where e is the edge $[0, 1]$ of I , and the sequence $j_1, \dots, j_p$ is increasing from 0 to p , omitting only i . Thus there are precisely $p + 1$ nondegenerate $p + 1$ simplices. Since $e = [0, 1]$, notice that all of the degeneracy maps before the “gap” at i must adjoin another 0 and all of those after the “gap” adjoin more 1s. Thus we can also label these nondegenerate $p + 1$ simplices exactly by the $p + 1$ sequences of length $p + 2$ of the form $[0, \dots, 0, 1, \dots, 1]$ that must start with a 0 and end with a 1.

If this looks familiar, it’s because the standard way to triangulate the product prism $\Delta^p \times I$ when studying simplicial homology theory is by the $p + 1$ simplices $[0, \dots, k, k', \dots, p']$, where the unprimed numbers represent vertices in $\Delta^p \times 0$ and the primed numbers represent vertices in $\Delta^p \times 1$. The simplex $[0, \dots, k, k', \dots, p']$ corresponds to $k + 1$ zeros and $p - k + 1$ ones. See Figure 20.




Figure 20: The realization of $|\Delta^2 \times \Delta^1|$ with nondegenerate 3-simplices $[0, 1, 2, 2']$, $[0, 1, 1', 2']$, and $[0, 0', 1', 2']$

For our upcoming discussion of simplicial homotopy, it’s also worth looking at how these simplices are joined together along their boundaries. Let’s first look from the point of view of writing the $p + 1$ simplices of $\Delta^p \times I$ in the form $S_k = [0, \dots, k, k', \dots, p']$, where $0 \leq k \leq p$. If $i < k$, then $d_i s = [0, \dots, i - 1, i + 1, \dots, k, k', \dots, p']$. But this can be thought of as a p

simplex of $[0, \dots, i-1, i+1, \dots, p] \times I$ and so is part of the boundary $\partial\Delta^p \times I$. Similar considerations hold if $k > i+1$. The interesting “interior cases” are

$$\begin{aligned} d_k S_k &= [0, \dots, k-1, k', \dots, p'] \\ d_{k+1} S_k &= [0, \dots, k, (k+1)', \dots, p']. \end{aligned}$$

To understand the assembly of the prism $\Delta^p \times I$, notice that $d_k S_k = d_k S_{k-1}$ for $k > 0$ and $d_{k+1} S_k = d_{k+1} S_{k+1}$ for $k < p$. This tells us how to glue the $p+1$ simplices together to form $|\Delta^p \times I|$.

In our other notation, if we have $S_k = (s_k E_p, s_p \dots s_{k+1} s_{k-1} \dots s_0 e)$, then for $i < k$ we have, using the axioms,

$$d_i S_k = (s_{k-1} d_i E_p, s_{p-1} \dots s_{k+1} s_{k-1} \dots s_i (d_i s_i) s_{i-1} \dots s_0 e) = (s_{k-1} d_i E_p, s_{p-1} \dots s_k s_{k-2} \dots s_0 e).$$

Notice that we use the axioms to “pass d_i through,” converting each s_j to s_{j-1} along the way, until it “annihilates” with the original s_i (leaving the previous s_{i+1} converted to the new s_i). We wind up with a p -simplex that is recognizable as a p -simplex in $d_i E_p \times I$. Similarly, for $i > k+1$, we get $d_i S_k = (s_k d_{i-1} E_p, s_{p-1} \dots s_k s_{k-2} \dots s_0 e)$. The two “interior” cases correspond to $d_k S_k$ and $d_{k+1} S_k$:

$$\begin{aligned} d_k S_k &= (d_k s_k E_p, s_{p-1} \dots s_{k+1} s_{k-2} \dots s_0 e) = (E_p, s_{p-1} \dots s_{k+1} s_{k-2} \dots s_0 e) \\ d_{k+1} S_k &= (d_{k+1} s_k E_p, s_{p-1} \dots s_{k+2} s_{k-1} \dots s_0 e) = (E_p, s_{p-1} \dots s_{k+2} s_{k-1} \dots s_0 e). \end{aligned}$$

These are not in $\partial\Delta^p \times I$. However, we do again see that $d_k S_k = d_k S_{k-1}$ for $k > 0$ and $d_{k+1} S_k = d_{k+1} S_{k+1}$ for $k < p$.

6 Simplicial objects in other categories

Before moving on to discuss simplicial homotopy, we pause to note that the categorical definition of simplicial sets suggests a sweeping generalization.

Definition 6.1. Let $\mathbf{Cat}$ be a category. A *simplicial object in $\mathbf{Cat}$* is a contravariant functor $X : \Delta \rightarrow \mathbf{Cat}$ (equivalent, a covariant functor $X : \Delta^{op} \rightarrow \mathbf{Cat}$). A morphism of simplicial objects in $\mathbf{Cat}$ is a natural transformation of such functors.

Another common notation, when $\mathbf{Cat}$ is a familiar category with objects of a given type, is to refer to a simplicial object in $\mathbf{Cat}$ as a simplicial [insert type of object]. In other words, when $\mathbf{Cat}$ is the category of groups and group homomorphisms, we speak of simplicial groups. This is consistent with referring to a simplicial object in the category $\mathbf{Set}$ as a simplicial set. One also commonly encounters simplicial R -modules, simplicial spaces, and even simplicial categories!

Example 6.2. Let’s unwind the definition in the case of simplicial groups. By definition, a simplicial group $\mathcal{G}$ consists of a sequence of groups $\mathcal{G}_n$ and collections of group homomorphisms $d_i : \mathcal{G}_n \rightarrow \mathcal{G}_{n-1}$ and $s_i : \mathcal{G}_n \rightarrow \mathcal{G}_{n+1}$, $0 \leq i \leq n$, that satisfy the axioms (1).

At this point, unfortunately, trying to picture group elements as simplices breaks down a little bit since there is so much extra structure around (what does it mean geometrically to multiply two simplices?). Nonetheless, it is still helpful to refer mentally to the category Δ , in which we can visualize each simplex $[n]$ as representing a group and picture movement toward each $n - 1$ face as representing a different group homomorphism to the group represented by $[n - 1]$. See Figure 21.




Figure 21: A pictorial representation of a 2-simplex of a simplicial group with arrows representing the face morphisms from dimension 2 to dimension 1 and from dimension 1 to dimension 0

Example 6.3. Suppose X is a simplicial set. Then we can form the simplicial group $C_*(X)$ with $(C_*X)_n = C_n(X)$ defined to be the free abelian group generated by the elements of X_n with d_i in $C_*(X)$ taken to be the linear extensions of the face maps d_i of X . We can also form the total face map $d = \sum_{i=0}^n (-1)^i d_i : C_n(X) \rightarrow C_{n-1}(X)$ and then define the homology $H_*(X)$ as the homology of this chain complex.

If $X = \mathcal{S}(Y)$, the singular set as defined in Example 3.7, then we have $H_*(X) = H_*(Y)$, the singular homology of the space Y .

Example 6.4. Here's an example of a simplicial group that is important in the theory of group cohomology. Let G be a group, and let BG be the simplicial group defined as follows. Let $BG_n = G^{\times n}$, the product of G with itself n times. $G^{\times 0}$ is just the trivial group $\{e\}$. For an element $(g_1, \dots, g_n) \in BG_n$, let

$$\begin{aligned} d_0(g_1, \dots, g_n) &= (g_1, \dots, g_n) \\ d_i(g_1, \dots, g_n) &= (g_1, \dots, g_i g_{i+1}, \dots, g_n) \quad \text{if } 0 < i < n \\ d_n(g_1, \dots, g_n) &= (g_1, \dots, g_{n-1}) \\ s_i(g_1, \dots, g_n) &= (g_1, \dots, g_i, e, g_{i+1}, \dots, g_n). \end{aligned}$$

The reader can check that this defines a simplicial group. The realization of the underlying simplicial set turns out to be the classifying space of the group G . For more on this simplicial group and its uses, the reader may consult [21, Chapter 8].

7 Kan complexes

One of the goals of the development of simplicial sets (and other simplicial objects) was to find a combinatorial way to study homotopy theory, just as simplicial homology theory allows us to derive invariants of simplicial complexes in a purely combinatorial manner (at least in principle). Unfortunately, it turns out that not all simplicial sets are created equal as regards their usefulness toward this goal. The underlying reason turns out to be (once again, at least in principle) related to the reason that homotopy theorists prefer to work with CW complexes and not arbitrary topological spaces. Pairs of CW complexes satisfy the homotopy extension property, i.e. inclusions of subcomplexes are cofibrations (see, e.g., [4]). The condition we need to impose on simplicial sets to make them appropriate for the study of homotopy is similarly an extension condition. When seen through sufficiently advanced lenses, such as from the viewpoint presented in [8], the extension condition on simplicial sets and the homotopy extension property in topology are essentially equivalent.

As with much else in the theory of simplicial sets, the extension condition comes from a fairly straightforward idea that is often completely obfuscated in the formal definition.

To explain the idea, we first need the following definition.

Definition 7.1. As a simplicial complex, the k th horn $|\Lambda_k^n|$ on the n -simplex $|\Delta^n|$ is the subcomplex of $|\Delta^n|$ obtained by removing the interior of $|\Delta^n|$ and the interior of the face $d_k \Delta^n$. See Figure 22. We let Λ_k^n refer to the associated simplicial set. This simplicial set consists of simplices $[i_0, \dots, i_m]$ with $0 \leq i_0 \leq \dots \leq i_m \leq n$ such that 1) not all numbers $0, \dots, n$ are represented (this would be the top face or a degeneracy thereof) and 2) we never have all numbers except k represented (this would be the missing $n - 1$ face or a degeneracy thereof).




Figure 22: The three horns on $|\Delta^2|$

The extension condition, also known as the Kan condition (after Daniel Kan), says that whenever we see a horn on an n -simplex within a simplicial set, the rest of the simplex is there, too. Here's an elegant way to say this:

Definition 7.2. The simplicial object X satisfies the *extension condition* or *Kan condition* if any morphism of simplicial sets $\Lambda_k^n \rightarrow X$ can be extended to a simplicial morphism $\Delta^n \rightarrow X$.

Such an X is often called a *Kan complex* or, in more modern language, is referred to as being *fibrant* (note the risk of confusion here between simplicial sets and simplicial complexes).

We next present an equivalent formulation that is often used. This version has its advantages from the point of view of conciseness of combinatorial information, but it is much less conceptual.

Definition 7.3 (Alternate version of the Kan condition). The simplicial set X satisfies the *Kan condition* if for any collection of $(n-1)$ -simplices $x_0, \dots, x_{k-1}, x_{k+1}, \dots, x_n$ in X such that $d_i x_j = d_{j-1} x_i$ for any $i < j$ with $i \neq k$ and $j \neq k$, there is an n -simplex x in X such that $d_i x = x_i$ for all $i \neq k$.

The condition on the simplices x_i of the alternative definition glues them together to form the horn Λ_k^n , possibly with degenerate faces, within X , and the definition says that we can extend this horn to a (possibly degenerate) n -simplex in X .

Example 7.4. Not even the standard simplices Δ^n , $n > 0$, satisfy the Kan condition! Let $\Delta^1 = [0, 1]$ be the standard 1-simplex, and consider the horn Λ_0^2 , which consists of the edges $[0, 2]$ and $[0, 1]$ of Δ^2 , along with their degeneracies. Now consider the simplicial morphism that takes $[0, 2] \in \Lambda_0^2$ to $[0, 0] \in \Delta^1$ and $[0, 1] \in \Lambda_0^2$ to $[0, 1] \in \Delta^1$. There is a unique such simplicial map since we've specified what happens on all the nondegenerate simplices of Λ_0^2 . Notice that this is perfectly well-defined as a simplicial map since all functions on all simplices are order preserving. However, this cannot be extended to a map $\Delta^2 \rightarrow \Delta^1$ since we have already prescribed that $0 \rightarrow 0$, $1 \rightarrow 1$, and $2 \rightarrow 0$, which is clearly not order preserving on Δ^2 .

Example 7.5. It is easy to check that Δ^0 does satisfy the Kan condition.

Example 7.6. Given a topological space Y , the simplicial set $\mathcal{S}(Y)$ *does* satisfy the Kan extension condition. It is actually fairly straightforward to see this. Consider any morphism of simplicial sets $f : \Lambda_k^n \rightarrow \mathcal{S}(Y)$. This is the same as specifying for each $n-1$ face, $d_i \Delta^n$, $i \neq k$, of Δ^n a singular simplex $\sigma_i : |\Delta^{n-1}| \rightarrow Y$. Every other simplex of Λ_k^n is a face or a degeneracy of a face of one of these $n-1$ simplices, and so the rest of the map f is determined by this data. Furthermore, the compatibility conditions coming from the simplicial set axioms ensure that the topological maps σ_i piece together to yield, collectively, a continuous function $f : |\Lambda_k^n| \rightarrow Y$. It is now a simple matter to extend this function to all of $|\Delta^n|$: let $\pi : |\Delta^n| \rightarrow |\Lambda_k^n|$ be any continuous retract (which certainly exists: $(|\Delta^n|, |\Lambda_k^n|)$ is homeomorphic to $(I^{n-1} \times I, I^{n-1} \times 0)$), and define $\sigma = f\pi : |\Delta^n| \rightarrow Y$. This is a singular n -simplex whose faces $d_i \sigma$, $i \neq k$, are precisely the singular simplices σ_i . Thus this determines the desired extension. See Figure 23.

Example 7.7. Any simplicial group is also, by neglect of structure, a simplicial set. All such simplicial sets arising from simplicial groups satisfy the Kan condition. The proof is not difficult, but I don't know of a version that is particularly illuminating. Since we will not have much further use for this fact in these notes, we refer the reader to [13, Theorem 2.2] for a proof.




Figure 23: A demonstration that the singular set satisfies the Kan condition

8 Simplicial homotopy

In this section we begin to look at the homotopy properties of simplicial sets. This is one of the key reasons that the theory of simplicial sets exists - to allow us to turn homotopy theoretic problems, at least in principle, into combinatorial problems by studying the homotopy groups of simplicial sets instead of those of topological spaces. In order to get started with simplicial homotopy, it is necessary to restrict attention to simplicial sets satisfying the Kan condition. This is not as large a handicap as it first appears, however, since we have already seen that, given a topological space Y , the singular set $\mathcal{S}(Y)$ satisfies the Kan condition, and eventually, we will see that $\mathcal{S}(Y)$ constitutes an appropriate combinatorial stand-in for Y .

As we proceed, the reader should bear in mind the extent to which many of the ideas and definitions mirror those in topological homotopy theory. This may prove a comfort (or cause serious worry!) at those junctures where the mirror appears somewhat warped by the combinatorial complexity of the simplicial version.

We begin, naturally enough, with π_0 , corresponding to the homotopy relationship between maps of points. This is a quite tractable warm-up for what is to follow.

8.1 Paths and path components

As in topology, when talking about homotopy, we will let I stand for the simplicial set $\Delta^1 = [0, 1]$. As a simplicial set, I has the nondegenerate 1-simplex $[0, 1]$, the nondegenerate 0-simplices $[0]$ and $[1]$, and all other simplices are degenerate. Each simplex has the form $[0, \dots, 0, 1, \dots, 1]$ (possibly with no 0s or no 1s).

Definition 8.1. A *path* in a simplicial set X is a simplicial morphism $p : I \rightarrow X$. Equivalently, a path in X is a 1-simplex $p \in X_1$. If p is a path in X , $d_1 p = p[0]$ is called the *initial point* of the path and $d_0 p = p[1]$ is called the *final point* or *terminal point*.

Definition 8.2. Two 0-simplices a and b of the simplicial set X are said to be *in the same path component* of X if there is a path p with initial point a and final point b .

Already this definition appears slightly odd if you're used to working with simplicial complexes. In a connected simplicial complex, one might have to traverse several edges to link two vertices. Here we require it to be done all with one edge. Furthermore, we would expect “being in the same path component” to be an equivalence relation. This is not at all clear, say, in an ordered simplicial complex in which we can have $a < b$ or $b < a$ but not both. What rescues this definition is the Kan condition.

Theorem 8.3. *If X is a Kan complex, then “being in the same path component” is an equivalence relation.*

Proof. We will go through the proof in detail as it is very illuminating of how to think geometrically about simplicial sets.

Reflexivity. This one is easy: for any vertex a , $s_0[a]$ is a path from a to a .

Transitivity. Consider $\Delta^2 = [0, 1, 2]$. If p_1 is a path from a to b and p_2 is a path from b to c , then let $f : \Lambda_1^2 \rightarrow X$ take $[0, 1]$ to p_1 and $[1, 2]$ to p_2 . See Figure 24. The Kan condition lets us extend f to $\bar{f} : \Delta^2 \rightarrow X$, and $\bar{f}[0, 2]$ gives us a path from a to c .




Figure 24: The transitivity relation on path connectedness via the Kan condition

Symmetry. This is only slightly more tricky than the transitivity condition. See Figure 25. Let p be a path in X from a to b . We need a path the other way. Think of p as the $[0, 1]$ side of Δ^2 . Let the $[0, 2]$ side of Δ^2 represent $s_0[a]$, which must exist since X is a simplicial set. Notice that $d_0 s_0[a] = d_1 s_0[a] = [a]$. At this point, we can label the three vertices $[0, 1, 2]$ of Δ^2 as $[a, b, a]$, and we have a simplicial map on Λ_0^2 taking $[0, 1]$ to p and $[0, 2]$ to $s_0[a]$. The Kan condition tells us that this map can be extended to all of Δ^2 and $[1, 2]$ gets taken to a path p from b to a . $\square$

Notice how important the Kan condition is here.

Since we have demonstrated that being in the same path component is an equivalence relation, we have equivalence classes.

Definition 8.4. We denote the set of path components of X (i.e. the equivalence classes of vertices of X under the relation of being in the same path component) by $\pi_0(X)$.

So far, this is comfortingly familiar.




Figure 25: The symmetry relation on path connectedness

8.2 Homotopies of maps

There are at least two classical versions of the definition of simplicial homotopy, and at least two more modern versions for which we refer the reader to [8]. Of the two classical versions, one has the expected form for a homotopy $H : X \times I \rightarrow Y$. The other is more closely related to the homotopies we see in chain complexes $\widehat{H} : X_n \rightarrow Y_{n+1}$. We will look at both of these and see how they are related.

Perhaps the most natural definition of simplicial homotopy looks something like this:

Definition 8.5 (Simplicial homotopy 1). Two simplicial maps $f, g : X \rightarrow Y$ are *homotopic* if there is a simplicial map $H : X \times I \rightarrow Y$ such that $H|_{X \times 0} = g$ and $H|_{X \times 1} = f$ (i.e., if $g = H \circ i_0$ and $f = H \circ i_1$, where i_0, i_1 are the evident simplicial inclusion maps $i_0 : X \times [0] \hookrightarrow X \times I$ and $i_1 : X \times [1] \hookrightarrow X \times I$).

Unfortunately, here is the definition of simplicial homotopy one finds quite often in the literature:

Definition 8.6 (Simplicial homotopy 2). Two simplicial maps $f, g : X \rightarrow Y$ are *homotopic* if for each p there exist functions $h_i = h_i^p : X_p \rightarrow Y_{p+1}$ for each i , $0 \leq i \leq p$, such that

1.

$$\begin{aligned} d_0 h_0 &= f \\ d_{p+1} h_p &= g \end{aligned}$$

2.

$$\begin{aligned} d_i h_j &= h_{j-1} d_i & \text{if } i < j \\ d_{j+1} h_{j+1} &= d_{j+1} h_j \\ d_i h_j &= h_j d_{i-1} & \text{if } i > j + 1 \end{aligned}$$

3.

$$\begin{aligned} s_i h_j &= h_{j+1} s_i & \text{if } i \leq j \\ s_i h_j &= h_j s_{i-1} & \text{if } i > j. \end{aligned}$$

It will take some doing to see how these two definitions are related. This was one of the initial motivations for writing this exposition!

As usual, we will consider the universal example, $X = \Delta^p$, since once we understand how a homotopy works on a single simplex, we will also understand what happens along its faces and degeneracies, and everything else is determined by how the simplices are glued together.

The key here is to recall Example 5.5 of Section 5, in which we showed how the prism $|\Delta^p \times I|$ is decomposed into simplices. In particular, it consists of $p + 1$ nondegenerate $p + 1$ simplices that we labeled $S_k \in (\Delta^p \times I)_{p+1}$, $0 \leq k \leq p$. Suppose now that we have a homotopy $H : \Delta^p \times I \rightarrow Y$ from f to g . Everything is determined by what H does to the S_k , since every other nondegenerate simplex in $\Delta^p \times I$ is a face of one of these simplices. All other simplices in $\Delta^p \times I$ are degenerate, and so their images are determined by the images of the simplices of which they are degeneracies.

How does this relate to the combinatorial Definition 8.6? Let us denote the unique nondegenerate p -simplex of Δ_p by E_p . In this version, there are $p+1$ functions $h_i : E_p \rightarrow Y_{p+1}$. Each of the $p + 1$ functions h_i assigns to E_p a $p + 1$ simplex of Y . Collectively, these give us the image of the prism over E_p in Y .

To see this, we use the notation $S_k = [0, \dots, k, k', \dots, p]$, $0 \leq k \leq p$, for the $p+1$ simplices of the prism $\Delta^p \times I$ (see Example 5.5). Given $H : \Delta^p \times I \rightarrow Y$, let $h_k(E_p)$ correspond to the image $H(S_k)$ in Y . Now let's look at the conditions in Definition 8.6 and see what they mean.

Starting with the first conditions, $d_0 h_0(E_p) = d_0 H(S_0) = H(d_0 S_0) = H(d_0[0, 0', \dots, p']) = H([0', \dots, p']) = H \circ i_1(E_p) = f(E_p)$, using the first definition of homotopy. Similarly, $d_{p+1} h_p(E_p) = H([0, \dots, p]) = H \circ i_0(E_p) = g(E_p)$. So these conditions assure that the ends of the prism really are controlled by the maps f and g .

The first and third equations of the second set of conditions mirror the observations made in Example 5.5 that most of the boundaries of the $p + 1$ simplices of the prism $\Delta^p \times I$ are themselves simplices of the prisms built on the boundary faces of Δ^p . So these equations ensure that these faces of the $h_i(\Delta^p)$ are compatible with the actions of the homotopy maps h_i^j of lower dimensions $j < p$ on the faces of Δ^p . The second equation is the condition that the neighboring simplices S_k and S_{k+1} share a boundary. We invite the reader to glean these combinatorial details from the calculations in Example 5.5.

The third set of equations can also be obtained in a fairly straightforward manner by working with the S_k . For example, we observe that for $i \leq j$, $s_i S_j = [0, \dots, i, i, \dots, j, j', \dots, p']$, which is also the $(j + 1)$ st prism simplex on the degenerate simplex $[0, \dots, i, i, \dots, j - 1, j, j + 1, \dots, p]$. In other words, the i th degeneracy of the j th prism $p + 1$ simplex over Δ^p is the $j + 1$ st prism simplex over the i th degeneracy of Δ^p . The geometric idea is a bit less obvious than in the preceding paragraphs, but really this is just the condition that the way the homotopy acts on degenerate simplices is determined by how it acts on the simplices of which they are degeneracies.

Having described how the combinatorial conditions of the second definition correspond to the more geometric ideas of the first definition, we now leave it to the interested reader to verify the complete equivalence of the two definitions, in particular to verify that the data given by all the h_i^j is enough to reconstruct H .

We would like for homotopy to be an equivalence relation, but this will not hold in general. For example, in our discussion of path connectedness, which we see in the current language corresponds directly to homotopies of maps $\Delta^0 \rightarrow X$, we saw that path connectedness is not always an equivalence relation. However, the discussion of path connectedness might lead one to suspect that we will be safe in the world of Kan complexes, and this is so.

Theorem 8.7. *Homotopy of maps $X \rightarrow Y$ is an equivalence relation if Y is a Kan complex. If f and g are homotopic, we denote that by $f \sim g$.*

We invite the reader to prove this by extending the argument given for path connectedness.

It is also fairly straightforward to verify other expected elementary fact about homotopy; for instance if $f \sim f'$, then $fg \sim f'g$ and $gf \sim gf'$. Also, homotopic maps induce the same homomorphisms on homology groups (see Section 3 - this follows as for the usual proof in singular homology theory by using the triangulation of the homotopy prism; see, e.g. [14]).

8.3 Relative homotopy

The notions of subcomplexes and relative homotopy offer no surprises, but we record the definitions for clarity.

Definition 8.8. If X is a simplicial set, then A is a simplicial subset of X , denoted $A < X$, if A itself is a simplicial set such that $A_n \subset X_n$ for all n and the face and degeneracy maps of A agree with those from X . The pair of simplicial sets is often denoted (X, A) . Simplicial maps of pairs $(X, A) \rightarrow (Y, B)$ are simplicial maps $X \rightarrow Y$ that such that the image of A is contained in B .

Definition 8.9. If (X, A) are a simplicial set and simplicial subset and both X and A satisfy the Kan condition, then (X, A) is called a Kan pair.

Example 8.10. An important example of a simplicial subset of a simplicial set X is a *basepoint* for X , consisting of an element of X_0 and all of its degeneracies. We will denote basepoints by $*$. Notice that $*$ is isomorphic as a simplicial set to Δ^0 and can be considered as an image $\Delta^0 \rightarrow X$ of a simplicial map. Since Δ^0 is a Kan complex, $(X, *)$ will be a Kan pair if X is Kan.

Example 8.11. Note that it is not automatic that a subcomplex of a Kan complex be Kan. For instance, we know from Example 7.4 that the simplex Δ^1 is not a Kan complex. We also know that the singular set $\mathcal{S}(|\Delta^1|)$ on the space $|\Delta^1|$ is a Kan complex, by Example 7.6. But the former is a subcomplex of the latter, realized by the singular simplices that represent $|\Delta^1|$ as a simplicial complex. Namely, Δ^1 corresponds to the subcomplex of $\mathcal{S}(|\Delta^1|)$ generated by the singular 0-simplices $\sigma_0 : |\Delta^0| \rightarrow [0]$ and $\sigma_1 : |\Delta^0| \rightarrow [1]$, by the singular 1-simplex $\text{id} : |\Delta^1| \rightarrow |\Delta^1|$, and by their degeneracies.

Definition 8.12. A homotopy $H : X \times I \rightarrow Y$ is a homotopy *rel* A if the restriction of H to $A \times I$ can be factored as $H|_{A \times I} = g\pi_1 : A \times I \rightarrow Y$, where g is a simplicial map $A \rightarrow Y$ and π_1 is the projection $A \times I \rightarrow A$. If Y is Kan, then homotopy *rel* A is an equivalence relation.

While considering simplicial pairs, there is another crucial theorem we should mention: the homotopy extension theorem for simplicial maps to Kan complexes:

Theorem 8.13 (Homotopy extension theorem.). *Let (X, A) be a pair of simplicial sets and Y a Kan complex. Suppose there is a simplicial map $f : X \rightarrow Y$ and a simplicial homotopy $H : A \times I \rightarrow Y$ such that $H|_{A \times 0} = f|_A$. Then there exists an extension $F : X \times I \rightarrow Y$ such that $F|_{A \times I} = H$ and $F|_{X \times 0} = f$.*

Unfortunately, the proofs I know would all take us too far afield, so we refer the reader to [13, Chapter 1, Appendix A] for a combinatorial treatment or [8, Section I.4] for a more modern treatment.

9 $\pi_n(\mathbf{X}, *)$

In this section, we will discuss the homotopy groups of Kan complexes. This section is a bit more technical than the preceding ones, as we here need some theorems and not just definitions. This section should serve as good technical practice for the reader preparing to go on to read further material on simplicial objects.

Given a simplicial set X , there are at least three ways to define $\pi_n(X, *)$:

1. One can define these groups directly as homotopy classes of maps of spheres.
2. There is a purely combinatorial definition.
3. As in algebraic topology, one can first define appropriate iterated simplicial loop spaces $\Omega^n(X)$ and define $\pi_n(X) = \pi_0(\Omega^n(X))$.

We will focus on the relationship between the first two of these, referring the interested reader to [13] for the third approach.

The definition of $\pi_n(X, *)$ in terms of spheres is straightforward once we decide what a sphere is. Example 4.7 teaches us that there is more than one reasonable definition, or at least more than one simplicial set whose realization is a sphere. In fact, we will see that both versions treated in that example are acceptable.

Definition 9.1 (First definition of π_n). Given a Kan complex with basepoint $(X, *)$, define $\pi_n(X, *)$, $n > 0$, to be the set of homotopy equivalence classes of maps $(\partial\Delta^{n+1}, *) \rightarrow (X, *)$. Here, we take for the basepoint of $\partial\Delta^{n+1}$ the simplicial subset of Δ^{n+1} generated by the vertex $[0]$, and all homotopies are relative to the basepoint.

The requirement that X be Kan is necessary for homotopy to be an equivalence relation. Of course, we want $\pi_n(X, *)$ to be a group, but this will have to wait a moment. Let's first work toward the more combinatorial definition. This takes a little bit of preliminary effort.

Definition 9.2. We say that two n -simplices $x, x' \in X_n$ are homotopic if

1. $d_i x = d_i x'$ for $0 \leq i \leq n$, and

2. there exists a simplex $y \in X_{n+1}$ such that

- (a) $d_n y = x$,
- (b) $d_{n+1} y = x'$, and
- (c) $d_i y = s_{n-1} d_i x = s_{n-1} d_i x'$, $0 \leq i \leq n-1$.

The idea here is that x and x' have the same boundaries and that y provides the homotopy between them, rel boundary, by letting x and x' be two of the faces of y , while the rest of the faces of y degenerate to the most straightforward degeneracies of the boundaries of x and y . See Figure 26.




Figure 26: Above: a homotopy of 1-simplices. Below: a homotopy of 2-simplices. The picture in the bottom right depicts two 2-simplices glued together along their boundaries.

It can be shown directly that homotopy of simplices is an equivalence relation if X is a Kan complex. The argument is a generalization of the one showing that path connectedness is an equivalence relation. Again the idea is to arrange a simplex so that the pieces we know fall on certain faces of horns and the pieces we'd like to show exist fall on the missing faces. Then these relations must exist due to the Kan extension condition. We refer the interested reader to [11, Section I.3].

Definition 9.3 (Second definition of π_n). Given a Kan complex with basepoint $(X, *)$, we can also define $\pi_n(X, *)$, $n > 0$, as the set of equivalence classes of simplices $x \in X$ with $d_i x \in *$ for all i , $0 \leq i \leq n$, up to homotopy of simplices.

This version of the homotopy groups corresponds more closely to our second version of the sphere in Example 4.7. Recall that, as a simplicial set, this version of the sphere S^n had only two nondegenerate simplices: one in dimension n and one in dimension 0. An n simplex of X all of whose faces live in $*$ can be thought of as the image of that simplicial version of S^n in X . Thus this definition of $\pi_n(X, *)$ also makes some geometric sense. However, there

are some obvious questions, such as: Why do the first and second definition of π_n agree? And where is the group structure we expect?

To answer the first question, we need a series of lemmas:

Lemma 9.4. *If X is Kan and $d_i x = d_i x'$ for all i , we obtain the same equivalence relation as in Definition 9.2 if we instead require that $d_r y = x$, $d_{r+1} y = x'$ for some $0 \leq r \leq n$, and $d_i y = d_i s_r x = d_i s_r x'$ for $i \neq r, r+1$.*

Proof. We refer the reader to [11] for the full proof, which is contained within Lemma 5.5 there. The idea is to show that the case of the definition using $r, r+1$ is equivalent to the version with $r+1, r+2$ for each relevant r . This is done using an extension argument by which one creates an $n+2$ simplex which has the two desired homotopies on two of the sides. We illustrated a low-dimensional case in Figure 27: Suppose that x, x' are 1-simplices and that we have a y with $d_0 y = x$, $d_1 y = x'$. We want to find a z with $d_1 z = x$, $d_2 z = x'$. We form the horn Λ_0^2 , shown flattened on the right of Figure 27. We embed y as $[0, 1, 3]$ (note that this maintains its orientation simplicially despite the oddities of the drawing). We let the other sides of the horn be appropriate degeneracies of x' . Notice that there is no trouble embedding this horn in X extending $y \in X$. Now the Kan condition assures us that we can extend this embedding to all of Δ^3 , including the remaining face $[1, 2, 3]$. We can check that this last face can be taken as the desired z (be careful to notice that $d_1[1, 2, 3] = [1, 3]$ and $d_2[1, 2, 3] = [1, 2]$).

The idea in higher dimensions is precisely the same; the extra faces of the horn that exist in higher dimensions contain other degeneracies of faces of x - see [11, Lemma 5.5], [3, Proposition 1.19]. $\square$




Figure 27: Shifting indices in the homotopy relation. Here $*$ represents $d_2 s_0 x = d_2 s_0 x'$, which is a degenerate 1-simplex, both vertices being the first vertex of x , which is also the first vertex of x' .

Lemma 9.5. *If X is Kan, two n -simplices $x, x' \in X$ are homotopic in the sense of Definition 9.2 if and only if the inclusion maps $f : \Delta^n \rightarrow X$ and $f' : \Delta^n \rightarrow X$ that represent x and x' are homotopic rel boundary as maps.*

Proof. Of course to say that f represents x means that f takes the nondegenerate n -simplex E_n of Δ^n to $x \in X$.

One direction of the argument is fairly straightforward. In order to show that f and f' are homotopic, it suffices to find a chain of $n+1$ simplices of dimension $n+1$, representing the

images of non-degenerate simplices of the prism $\Delta^n \times I$, such that the “top” and “bottom” faces of the first and last simplex represents x and x' . But if we know that x and x' are homotopic as simplices, we know there is one $n + 1$ simplex y connecting them with, say, $d_n y = x$, $d_{n+1} y = x'$, and $d_i y \in *$ for all other i . So now we just let y be the $n + 1$ st simplex $h_n(\Delta^n)$, and we let $h_i(\Delta^n) = s_i x$ for $0 \leq i \leq n$. In other words, we let the last nondegenerate simplex in $\Delta^n \times I$ do the work of the homotopy, and we just collapse all the rest into the face representing x . See Figure 28.




Figure 28: We label the vertices with the “prism notation” of Example 5.5. The bottom simplex y is a homotopy of the simplices x and x' . Adjoining the degenerate simplex $s_0 x$ shows how to obtain a model prism for the homotopy from x to x' as inclusion maps.

In the other direction, suppose we have an actual homotopy $\text{rel } *$ from x to x' thought of as inclusion maps. By definition, this gives us a prism $\Delta^n \times \Delta^1 \in X$ whose top is x and whose bottom is x' . We know from the discussion in Example 5.5 that each of the nondegenerate $n + 1$ simplices of the prism have two n -faces that are not in $\partial \Delta^n \times \Delta^1$, and the rest are in $\partial \Delta^n \times \Delta^0$, all of which goes to $*$ in X . Furthermore, it is not hard to check that the two faces n -faces not in $\partial \Delta^n \times \Delta^1$ are consecutive faces. In particular, using the notation of Example 5.5, these faces are $d_k S_k$ and $d_{k+1} S_k$. Thus by Lemma 9.4, each S_k is a homotopy between these two faces. Since the top and bottom faces of the prism are x and x' , we obtain a simplicial homotopy between x and x' using the transitivity of simplicial homotopy. $\square$

Thus, to show that our two definitions of $\pi_n(X, *)$ agree, it is only necessary to prove the following lemma, which is familiar in the context of algebraic topology. The proof is somewhat long, but we provide most of the details, as it is difficult to find a direct proof in the standard expositions.

Lemma 9.6. *If X is a Kan complex, there is a bijection between homotopy classes of maps $f : (\partial \Delta^{n+1}, *) \rightarrow (X, *)$ and homotopy classes of maps $g : (\Delta^n, \partial \Delta^n) \rightarrow (X, *)$.*

Proof. Given $g : (\Delta^n, \partial \Delta^n) \rightarrow (X, *)$, it is easy to construct an associated f by identifying Δ^n with $d_0 \Delta^{n+1}$. Then we let $f : (\partial \Delta^{n+1}, *) \rightarrow (X, *)$ be defined so that f is given by g on $d_0 \Delta^{n+1}$ and by the unique map to $*$ on each $d_i \Delta^{n+1}$, $i > 0$. It is also straightforward to see that any homotopy of $g \text{ rel } \partial \Delta^n$ determines a homotopy of $f \text{ rel } *$.

Conversely, suppose we are given $f : (\partial\Delta^{n+1}, [0]) \rightarrow (X, *)$. We show that f is homotopic to a function $\tilde{f}$ that takes Λ_0^{n+1} to $*$. Then we can let g be $f|_{d_0\Delta^{n+1}}$.

We first observe, as note in the proof of Lemma 9.5, that to construct a homotopy between two inclusions of k -simplices x and x' in X , it suffices to find a simplex y in X with $d_k y = x$, $d_{k+1} y = x'$ since this can be considered one of the blocks of a prism, and the rest of the prism can be filled up with degeneracies of x or x' .

Keeping this in mind, we proceed by induction with the following induction step: Suppose $f_{k-1} : \partial\Delta^{n+1} \rightarrow X$ is such that $f([0]) \in *$ and $f(z) \in *$ for all simplices $z \in \partial\Delta^{n+1}$ of dimension $\leq k-1$ such that $[0]$ is a vertex of z , then there is a homotopy from f_{k-1} to an f_k that takes all simplices up to dimension k having $[0]$ as a vertex to $*$. Furthermore, the homotopy can be performed rel the faces of dimension $\leq k-1$ having $[0]$ as a simplex.

Clearly we can take $f_0 = f$. So suppose we have constructed f_{k-1} for $k \geq 1$. We need only find the desired homotopy on the k simplices of Δ^{n+1} that have $[0]$ as a vertex, and then we can apply the homotopy extension theorem, Theorem 8.13. So let z be a k -simplex of Δ^{n+1} with 0 as a vertex. We know that $f_{k-1}(d_i z) \in *$ for $i \neq 0$. Now, consider the horn Λ_0^{k+1} , and note that we can map Λ_0^{k+1} into X such that the k -face corresponding to $d_{k+1}\Delta^{k+1}$ is $f_{k-1}(z)$ and such that all other k -faces are taken into $*$. Notice that this is possible precisely because $f_{k-1}(d_i z) \in *$ for $i \neq 0$. Now since X is a Kan complex, we can extend this horn to a $(k+1)$ -simplex y in X such that $d_{k+1}y = f_{k-1}(z)$ and $d_k y \in *$. As noted, this is enough to construct a homotopy on z from $f_{k-1}(z)$ to the unique map of z into $*$. In addition, this is a homotopy rel those faces of z that have $[0]$ as a simplex. Notice also that it is possible to find such homotopies for all such z independently and compatibly. In this way, we get a homotopy on the k -simplices of Δ^{n+1} having $[0]$ as a vertex from f_{k-1} to the map to $*$. Extending this homotopy by the homotopy extension theorem yields the desired homotopy to f_k .

Continuing inductively, we obtain a function $f_{n+1} : \partial\Delta^{n+1} \rightarrow X$ homotopic to f such that Λ_0^{n+1} is taken to $*$. Now we can define g to be the restriction of f_{n+1} to $d_0\Delta^{n+1}$.

If $f, f' : (\partial\Delta^{n+1}, [0]) \rightarrow (X, *)$ are homotopic rel $[0]$, then we can show that the resulting g and g' are homotopic by building a homotopy from the homotopy $H : \partial\Delta^{n+1} \times I \rightarrow X$ from f to f' to a homotopy $H_{k+1} : \partial\Delta^{n+1} \times I \rightarrow X$ such that $H_{k+1}(\Lambda_0^{n+1} \times I) \in *$ and that extends the homotopies built over f and f' as in the preceding paragraphs. Then $H_{k+1}|_{d_0\Delta^{n+1} \times I}$ will be a homotopy from g to g' . We leave the details to the reader. $\square$

Lemmas 9.5 and 9.6 together prove the following.

Proposition 9.7. *If X is a Kan complex, the definitions of $\pi_n(X, *)$ in Definitions 9.1 and 9.3 agree.*

The group structure. One benefit of the version of $\pi_n(X, *)$ given in Definition 9.3, compared to the perhaps more geometrically transparent Definition 9.1, is the ease of proving that $\pi_n(X, *)$ is a group and of describing the group operation.

Definition 9.8. Let x, y be two n -simplices, $n \geq 1$, in the Kan complex X such that $d_i x = d_i y \in *$ for all i . Let Λ_n^{n+1} be the horn of Δ^{n+1} in X such that the face corresponding

to $d_{n+1}\Delta^{n+1}$ is y , the face corresponding to $d_{n-1}\Delta^{n+1}$ is x , and the faces corresponding to all other sides of the horn are in $*$. Let z be an extension of the horn to Δ^{n+1} as guaranteed by the Kan condition. Then define xy as the homotopy class of $d_n z$ in $\pi_n(X, *)$. See Figure 29.




Figure 29: The product of x and y in $\pi_1(X, *)$ (above) or $\pi_2(X, *)$ (below).

It can be shown that the definition is independent of the choices made:

Proposition 9.9. *The product of Definition 9.8 yields a well-defined function $\pi_n(X, *) \times \pi_n(X, *) \rightarrow \pi_n(X, *)$.*

Proof. The proof is by various applications of the Kan extension condition. See [11, Lemma 4.2]. This would also be a good exercise for the reader. $\square$

The idea of the product on the simplicial $\pi_n(X, *)$ is not far from that for the product in the topological homotopy groups. First, suppose one has a map of the $(n+1)$ -ball D^{n+1} to a topological space X such that the equator of the boundary sphere S^n is mapped to the basepoint of X . Then the restrictions of the map to the upper and lower hemispheres of S^n determine elements of $\pi_n(X, *)$, and the map of all of D^{n+1} determines a homotopy between them. Secondly, recall that, roughly speaking, the product of two elements x, y in the topological $\pi_n(X, *)$ can be represented by a map of a sphere that agrees with x and y on two disjoint disks in S^n and takes the rest of S^n to the basepoint.

Definition 9.8 puts these ideas together. In the simplicial world, we can think of $d_n\Delta^{n+1}$ as being one hemisphere of $\partial\Delta^{n+1}$ and the rest of $\partial\Delta^{n+1}$ as the other hemisphere. Then in Definition 9.8, the $(n+1)$ -simplex z can be thought of as providing a homotopy between $d_n z$ and what is happening on the rest of ∂z (notice that, indeed, $\partial d_n z \in *$). But the rest of ∂z contains x and y on two separate faces and everything else goes to $*$, just as for the topological product.

Of course we expect $\pi_n(X, *)$ to be a group if $n > 0$.

Theorem 9.10. *With the product of Definition 9.8, $\pi_n(X, *)$ is a group.*

Proof. The constructions are pictured in Figure 30.




Figure 30: Above left: the identity $x* = x$. Above right: construction of the right and left inverses of x . Below: Associativity $(xy)z = x(yz)$.

The constant map $\Delta^n \rightarrow *$ (which we will also denote by $*$) is the identity element. Indeed, given $x \in X$ representing an element of $\pi_n(X, *)$, the $(n+1)$ -simplex $s_n x$ will have $d_{n+1}s_n x = d_n s_n x = x$, while for $i < n$, $d_i s_n x = s_{n-1} d_i x \in *$. This realizes $x = *x$. Similarly, consideration of $s_{n-1} x$ gives $x = x*$.

It is also easy to construct inverses: given $x \in X$ representing an element of $\pi_n(X, *)$, there is no problem mapping the horn Λ_{n+1}^{n+1} into X such that the face corresponding to $d_{n-1}\Delta^{n+1} = x$ and all other faces are in $*$. The Kan condition lets us extend this to a map of Δ^{n+1} into X and then the face corresponding to $d_{n+1}\Delta^{n+1}$ is a right inverse to x . Similarly, we can find a left inverse using Λ_{n-1}^{n+1} and putting x on the face corresponding to $d_{n+1}\Delta^{n+1}$.

Finally, we show that the group operation is associative, which takes a bit more work. Let x, y, z be simplices in X representing elements of $\pi_n(X, *)$. We choose $(n+1)$ -simplices w_{n-1} and w_{n+2} that respectively realize the products xy and yz , and we choose a simplex w_{n+1} realizing the product $(xy)z$, where xy is represented by $d_n w_{n-1}$. Now, we can find a horn Λ_n^{n+2} in X such that the faces corresponding to $d_i \Delta^{n+2}$ are the w_i for $i = n-1, n+1, n+2$ and $*$ otherwise. To see that this data is consistent to form the horn, we need to check the appropriate faces, most of which are in $*$, to see that they correspond. The only faces of Δ^{n+2} we don't need to check are those of the form $d_i d_n \Delta^{n+2}$ since $d_n \Delta^{n+2}$ isn't in the horn. By the simplicial axioms, these also correspond to the faces $d_{n-1} d_i \Delta^{n+2}$ for $i < n$ and $d_n d_{i+1} \Delta^{n+2}$ for $i \geq n$. This leaves the following faces to check: We have $d_n d_{n-1} \Delta_n^{n+2} = d_n w_{n-1} = xy = d_{n-1} w_{n+1} = d_{n-1} d_{n+1} \Delta_n^{n+2}$ and $d_{n+1} d_{n-1} \Delta_n^{n+2} = d_{n+1} w_{n-1} = y = d_{n-1} w_{n+2} = d_{n-1} d_{n+2} \Delta_n^{n+2}$. We also have $d_{n+1} d_{n+1} \Delta_n^{n+2} = d_{n+1} w_{n+1} = z = d_{n+1} w_{n+2} = d_{n+1} d_{n+2} \Delta_n^{n+2}$. All other sides in the proposed horn are in $*$, and so the data is consistent.

We can extend this horn to an $(n+2)$ -simplex u by the Kan condition. So now by definition of w_{n+1} , $(xy)z = d_n w_{n+1} = d_n d_{n+1} u$, which, using the axioms, is also equal to $d_n d_n u$. But this also represents the product of $d_{n-1} d_n u = d_{n-1} d_{n-1} u = d_{n-1} w_{n-1} = x$ with

$d_{n+1}d_nu = d_nd_{n+2}u = d_nw_{n+2} = yz$. So d_nd_nu also represents the product $x(yz)$, proving associativity. $\square$

Also as expected, $\pi_n(X, *)$ is an abelian group for $n \geq 2$, but this is a bit more difficult to prove. We refer the reader to [11, Proposition 4.4].

Relative homotopy groups. If $(X, A, *)$ is a Kan triple (meaning A is a Kan subcomplex of the Kan complex X and $*$ is a basepoint in A), there are also relative homotopy groups $\pi_n(X, A, *)$. Corresponding to our first definition of $\pi_n(X, *)$ and the topological notion of relative homotopy, we could define $\pi_n(X, A, *)$ to be relative homotopy classes of maps $(\Delta^n, \partial\Delta^n, [0]) \rightarrow (X, A, *)$, where the homotopies are required to keep the image of $\partial\Delta^n \times I$ in A and the image of $[0] \times I$ in $*$. For a version of $\pi_n(X, A, *)$ corresponding to our second definition of $\pi_n(X, *)$, we first need a relative notion of homotopy of simplices:

Definition 9.11. If A is a subcomplex of X , we say that two n -simplices $x, x' \in X_n$ are homotopic rel A if $d_ix = d_ix'$ for $1 \leq i \leq n$, d_0x is homotopic to d_0x' in A via an n -simplex y , and there exists a simplex $w \in X_{n+1}$ such that $d_0w = y$, $d_nw = x$, $d_{n+1}w = x'$, and $d_iw = s_{n-1}d_ix = s_{n-1}d_ix'$, $1 \leq i \leq n-1$.

This definition is very similar to that for homotopy of simplices except instead of requiring $d_0x = d_0x'$, we let d_0x and d_0x' be two simplices that are themselves homotopic in A , and the homotopy between x and x' , provided by w , contains within it the homotopy between d_0x and d_0x' .

Using this relative notion of homotopy, we can define $\pi_n(X, A, *)$.

Definition 9.12. Given a Kan triple $(X, A, *)$, we define $\pi_n(X, A, *)$, $n > 0$, as the set of equivalence classes of n -simplices $x \in X$ with $d_0x \in A$ and $d_ix \in *$ for all i , $1 \leq i \leq n$, up to relative homotopy of simplices.

$\pi_n(X, A, *)$ is also a group for $n \geq 2$ and an abelian group for $n \geq 3$. We will define the product; the proofs of well-definedness and that we have a group are analogous to that for $\pi_n(X, *)$.

Definition 9.13. Suppose x, y represent elements of $\pi_n(X, A, *)$, $n \geq 2$. Let z represent the product between d_0x and d_0y in $\pi_{n-1}(A, *)$. So $z \in A$ is such that $d_{n-2}z = d_0x$, $d_nz = d_0y$ and $d_{n-1}z$ represents $(d_0x)(d_0y)$. Now map the horn Λ_n^{n+1} into X such that the sides corresponding to $d_0\Delta^{n+1}$, $d_{n-1}\Delta^{n+1}$, and $d_{n+1}\Delta^{n+1}$ are z , x , and y , respectively, and all other faces go to $*$. One can check that this is consistent data. Then let w be an extension of the horn, which exists because X is Kan, and define xy to be d_nw .

An excellent exercise for the reader at this point would be to show that there is a long exact sequence

$$\rightarrow \pi_n(A, *) \rightarrow \pi_n(X, *) \rightarrow \pi_n(X, A, *) \rightarrow \pi_{n-1}(A, *) \rightarrow .$$




Figure 31: The product of two elements of $x, y \in \pi_2(X, A, *)$. The 1-simplex with endpoints 1 and 3 represents the product $(d_0x)(d_0y)$ in $\pi_1(A)$.

10 Concluding remarks

It is difficult to know where to end a survey of the type we have undertaken here. On the one hand, although we have included some material from its later chapters, we have not even covered the entire first chapter of May's textbook [11]! On the other hand, our goal has never been to provide a completely rigorous or comprehensive treatise on simplicial theory, but to provide the reader with an introduction to some of the most important elementary ideas while maintaining a bridge to the geometric pictures that the combinatorics are based upon. We hope that we have prepared the interested student to move on to the more standard texts on simplicial objects with some picture (literally) of what's going on there.

And what is going on there? Just about everything in topological homotopy theory and then some. Just a glance at the table of contents of [11] turns up many familiar concepts from homotopy theory: fibrations, fiber bundles, Postnikov systems, function spaces, Hurewicz theorems, Eilenberg-MacLane complexes, k -invariants, cup and cap products, the Serre spectral sequence... This is not surprising in light of the following theorem; we refer the reader to Curtis [3, Section 12], or to [8, Section I.11] for a modern proof.

Theorem 10.1. *The homotopy category of Kan complexes, consisting of Kan complexes and homotopy classes of maps between them, is equivalent to the category of CW complexes and homotopy classes of continuous maps.*

The functors that realize this equivalence are the realization functor of simplicial complexes and the singular set functor that assigns the singular set to a topological space. Thus this theorem is closely related to the adjointness theorem, Theorem 4.10. So this tells us that everything we have been doing in the simplicial realm is a reflection of ordinary homotopy theory. Yet, despite the geometric point of view we have been emphasizing here, simplicial theory is purely combinatorial and algebraic, accessible by discrete tools that may not be evident in pure topology. Thus, using simplicial theory, one can hope to study topological homotopy theory via these combinatorial tools. Furthermore, we touched upon how the combinatorial simplicial methods can be transported to other contexts, such as simplicial groups. They can also be abstracted to broader categorical settings, leading to the theory

of model categories and simplicial categories. We hope to have introduced enough of the background also to enable the reader to pursue these more modern approaches, such as found in [8], with some understanding of their original motivation in concrete homotopy theory.

We leave the reader with some bibliographical notes on the sources we have used.

Our primary sources were May's *Simplicial Objects in Algebraic Topology* [11] and Moore's lecture notes *Seminar on algebraic homotopy theory* [13]. May's book, first published in 1967, is the most comprehensive reference of its time, featuring a direct combinatorial approach. Moore's notes are from nearly a decade earlier, but they are perhaps a bit more accessible to the geometrically-minded reader; they take a different approach to homotopy groups, defining them as π_0 of simplicial loops spaces. Our primary modern source was *Simplicial Homotopy Theory* [8] by Goerss and Jardine. It starts off directly from the modern model category point of view, without much need for the combinatorial underpinnings (some knowledge of the combinatorial approach, however, will aid the reader). Despite the abstractness of the material, I found this book quite readable. The book *Calculus of Fractions and Homotopy Theory* [7] by Gabriel and Zisman, though contemporary with May's book, is something of a bridge between the classical combinatorics and some of the more current axiomatic ideas. We should also mention in this paragraph the long survey *Simplicial Homotopy Theory* [3] by Curtis. As one might expect, each of these sources contains somewhat different material and sometimes different approaches to the same material, thus it is well worth consulting each of them depending on the reader's interests in terms of both material and style.

Besides these longer expositions, short introductory chapters on simplicial theory can be found within many other textbooks and surveys. In particular, I know of sections on simplicial theory in Selick's *Introduction to Homotopy Theory* [19], Smirnov's *Simplicial and Operad Methods in Algebraic Topology* [20], and Weibel's *An Introduction to Homological Algebra* [21]. As one might expect, this last reference is a good source for applications of simplicial theory to homological algebra. There are also review sections on simplicial sets in Bousfield and Kan's *Homotopy Limits, Completions, and Localizations* [1] and in *Mixed Hodge Structures* [15] by Peters and Steenbrink. The breadth of topics covered by those titles alone should give the reader some impression of just how varied the applications of simplicial theory are.

References

- [1] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Mathematics, Vol. 304, Springer-Verlag, Berlin, 1972.
- [2] Glen Bredon, *Topology and geometry*, Springer-Verlag, New York, 1993.
- [3] Edward B. Curtis, *Simplicial homotopy theory*, *Advances in Math.* **6** (1971), 107–209.
- [4] James F. Davis and Paul Kirk, *Lecture notes in algebraic topology*, Graduate Studies in Mathematics, vol. 35, American Mathematical Society, Providence, RI, 2001.

- [5] Samuel Eilenberg and J. A. Zilber, *Semi-simplicial complexes and singular homology*, Ann. of Math. (2) **51** (1950), 499–513.
- [6] Yves Félix, Stephen Halperin, and Jean-Claude Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics, vol. 205, Springer-Verlag, New York, 2001.
- [7] P. Gabriel and M. Zisman, *Calculus of fractions and homotopy theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35, Springer-Verlag New York, Inc., New York, 1967.
- [8] Paul G. Goerss and John F. Jardine, *Simplicial homotopy theory*, Progress in Mathematics, vol. 174, Birkhauser Verlag, Basel, 1999.
- [9] Allen Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002.
- [10] J. P. May, *The geometry of iterated loop spaces*, Springer-Verlag, Berlin, 1972, Lectures Notes in Mathematics, Vol. 271.
- [11] J. Peter May, *Simplicial objects in algebraic topology*, University of Chicago Press, Chicago, IL, 1992.
- [12] John Milnor, *The geometric realization of a semi-simplicial complex*, Ann. of Math. (2) **65** (1957), 357–362.
- [13] J.C. Moore, *Seminar on algebraic homotopy theory*, Mimeographed notes - Princeton, 1956.
- [14] James R. Munkres, *Elements of algebraic topology*, Addison-Wesley, Reading, MA, 1984.
- [15] Chris A. M. Peters and Joseph H. M. Steenbrink, *Mixed Hodge structures*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 52, Springer-Verlag, Berlin, 2008.
- [16] Daniel Quillen, *Rational homotopy theory*, Ann. of Math. (2) **90** (1969), 205–295.
- [17] Daniel G. Quillen, *Homotopical algebra*, Lecture Notes in Mathematics, No. 43, Springer-Verlag, Berlin, 1967.
- [18] C. P. Rourke and B. J. Sanderson, Δ -sets. I. *Homotopy theory*, Quart. J. Math. Oxford Ser. (2) **22** (1971), 321–338.
- [19] Paul Selick, *Introduction to homotopy theory*, Fields Institute Monographs, vol. 9, American Mathematical Society, Providence, RI, 1997.
- [20] V. A. Smirnov, *Simplicial and operad methods in algebraic topology*, Translations of Mathematical Monographs, vol. 198, American Mathematical Society, Providence, RI, 2001, Translated from the Russian manuscript by G. L. Rybnikov.

- [21] Charles A. Weibel, *An introduction to homological algebra*, Cambridge studies in advanced mathematics, Cambridge University Press, Cambridge, 1994.

Some diagrams in this paper were typeset using the $\text{\TeX}$ commutative diagrams package by Paul Taylor.